



Metabolic Supremacy Fuels Tumor Aggressiveness in Renal Cancer

Reham Gholam¹, Muhammad Khalilzad²

Cite this article: Gholam R, Khalilzad M: Metabolic Supremacy Fuels Tumor Aggressiveness in Renal Cancer. *Ann Urol Oncol* 2025, 8(1): 40-49. <https://doi.org/10.32948/auo.2025.03.01>

Abstract

Renal cell carcinoma, with clear cell renal carcinoma (ccRCC) being the dominant form, is recognized as a malignancy driven by abnormal metabolic processes, with extensive alterations in glucose, lipid, and amino acid pathways. The loss of the Von Hippel-Lindau (VHL) gene in nearly 90% of ccRCC instances results in the accumulation of hypoxia-inducible factors (HIFs), producing a pseudo-hypoxic environment that promotes metabolic supremacy. This, in return, amplifies glucose uptake and directs energy production toward aerobic glycolysis, commonly referred to as the Warburg effect, even under conditions of good oxygen supply. Simultaneously, suppression of oxidative phosphorylation and heightened activity of the pentose phosphate pathway foster biosynthetic requirements and support an immunosuppressive environment. Dysregulated lipid metabolism, including elevated fatty acid synthesis, excessive cholesterol storage, and reduced β -oxidation, also contributes to disease aggressiveness. ccRCC cells also exhibit a pronounced reliance on glutamine, powering the tricarboxylic acid (TCA) cycle and preserving redox homeostasis, whereas altered tryptophan and arginine pathways facilitate immune escape. Overall, this metabolic supremacy fuels malignant growth, promote tumor aggressiveness and metastatic spread, and foster resistance to therapy. The pursuit of interventions targeting in this regard has been promising with HIF-2 α inhibitors, such as belzutifan, showing clinical benefit. Other emerging strategies focus on disrupting glycolysis, lipid biogenesis, and glutamine utilization in tackling metabolic supremacy in renal cancer. This comprehensive review delves into ccRCC's multifaceted metabolic landscape with focus on underlying pivotal molecular pathways, their implications in tumor aggressiveness, and the potential of innovative treatments targeting metabolic supremacy to limit tumor burden and improve patient outcomes in this malignancy.

Key words renal cancer, hypoxia, Warburg effect, lipid metabolism, amino acid metabolism

1. Department of Pharmacognosy, Daras Gah Mili Building, Islamic Azad University, ZIP area 11, Azarshahr Street, Karimkhan-e-Zand Avenue, Tehran 1477893780, Iran.

2. Department of Basic Medical Science, Block 15, Islamic Azad University, ZIP area 11, Azarshahr Street, Karimkhan-e-Zand Avenue, Tehran 1477893780, Iran.

Correspondence: Muhammad Khalilzad (Department of Basic Medical Science, Block 15, Islamic Azad University, ZIP area 11, Azarshahr Street, Karimkhan-e-Zand Avenue, Tehran 1477893780, Iran; Email: Khalilzadah888@yahoo.com).

Introduction

Cancer is driven by unregulated cell growth due to continuous proliferation leading to malignant transformation. A key enabler of this pathological expansion is the reprogramming of metabolic pathways, supporting increased energy demands and biosynthetic processes [1, 2]. Kidney cancer is especially relevant on a global scale, and was ranked as the 14th most commonly diagnosed cancer in 2020, accounting for 431,288 new cases and the 15th deadliest cancer with 179,368 death worldwide [3]. In the United States, nearly 79,000 new cases of kidney cancer were reported in 2022 alone, underscoring its clinical significance [4]. Clear cell renal carcinoma (ccRCC), arising from renal tubular epithelial cells, representing about 75% of all kidney cancer diagnoses [5, 6]. Although surgical procedures have advanced, ccRCC remains challenging to treat due to its aggressive behavior and metastatic potential, often necessitating adjunct therapies like targeted agents, immunotherapy combinations, or dual immunotherapy [7]. While these modalities have extended survival, their effectiveness is frequently limited by adverse effects and variability in patient response [8, 9].

First identified several decades ago, metabolic reprogramming has emerged as a primary focal point for new cancer treatments. It enables tumors to adapt biochemical pathways to satisfy escalated energy requirements and biosynthesis, thus presenting novel targets for intervention [10]. Early studies characterized the heightened glycolysis in cancer cells, known as the “Warburg effect”, as a reflection of inefficient energy generation [11]. However, modern evidence implicates oncogenic mutations in driving widespread metabolic alterations, influencing glucose uptake, lipid synthesis, and mitochondrial oxidative phosphorylation to advance tumorigenesis [12]. Even when oxygen is plentiful, cancer cells deplete local nutrient stores by favoring glycolysis over oxidative phosphorylation and establish an immunosuppressive environment that impairs T-cell function, thereby promoting tumor progression [13]. These metabolic abnormalities are particularly pronounced in ccRCC, with genetic disruptions triggering shifts in glucose metabolism, increased glutamine reliance, and mitochondrial dysfunction [14, 15]. The Warburg effect, exemplified by the bias toward rapid ATP generation while also producing building blocks vital for cellular proliferation, is a hallmark of ccRCC [16]. Hence, recognizing the relationship between metabolic dysregulation and ccRCC pathogenesis is essential for designing more effective treatments. Here, we explore the molecular mechanisms behind metabolic supremacy in ccRCC, focusing on dysfunctional hypoxic signaling, aberrant glucose metabolism, Warburg effect, dysregulated amino acid metabolism and elevated lipid metabolism as key culprits behind tumor aggressiveness. We also discuss the pharmacological interventions targeting these pathways, including agents that inhibit glycolysis, glutaminolysis, and fatty acid oxidation. By mapping the metabolic framework of ccRCC, this review aims to guide future research and therapeutic innovations intended to overcome tumor aggressive in renal cancer, critical to limit tumor burden and enhance patient survival in this malignancy.

Hypoxia signaling induces metabolic supremacy in renal cancer

ccRCC is often associated with alterations in the Von Hippel-Lindau (VHL) gene, which is associated with metabolic supremacy in up to 90% of cases [12]. Located on chromosome 3's short arm, VHL encodes the tumor suppressor protein pVHL [17]. Under normal conditions, pVHL facilitates the ubiquitination of proline-rich residues in the oxygen-dependent degradation domains of hypoxia-inducible factors (HIFs), directing them

toward proteasomal degradation [18, 19]. HIFs function as transcriptional regulators that enable cellular adaptation to low-oxygen environments [20]. Among the HIF isoforms, HIF-1 α is predominant in most tissues and cells, with its stability influenced by both oxygen availability and metabolic status [21, 22]. By contrast, under pathological hypoxia, HIF-2 α resists degradation, forms a complex with HIF-1 β , also referred to as the aromatic hydrocarbon receptor nuclear transporter (ARNT), and translocates into the nucleus to initiate gene transcription (**Figure 1**) [23]. In ccRCC, VHL mutations lead to the accumulation of HIF-2 α , creating a “pseudo-hypoxic” state [24]. This condition triggers metabolic reprogramming that fuels angiogenesis, epithelial-mesenchymal transition, tumor invasion, and metastasis [25]. The HIF pathway elevates the expression of key enzymes and transporters involved in glucose uptake and glycolysis, glucose transporter type 1 (GLUT1), phosphoglycerate kinase (PGK), lactate dehydrogenase (LDHA), pyruvate dehydrogenase kinase (PDK1), and hexokinase (HK) [26], while concurrently suppressing oxidative phosphorylation and the tricarboxylic acid (TCA) cycle [27]. Furthermore, ccRCC exhibits additional mutations in components of the PI3K-AKT-mTOR signaling cascade, including PTEN, TSC1/2, and PIK3CA [28, 29]. The proteins produced by TSC1 and TSC2 ordinarily act as an inhibitory complex for mTORC1 [30]. Once mTORC1 is activated, it suppresses the tumor suppressor 4EBP1, which boosts the expression of both HIF-1 and HIF-2, thereby amplifying the metabolic reprogramming that drives tumor development [31, 32]. Hence, the pseudo-hypoxia driven metabolic supremacy underscores the aggressive character of ccRCC and illustrates how genetic mutations and metabolic shifts intersect to propel tumor progression (**Figure 1**).

Aberrant glucose metabolism drives metabolic supremacy-driven tumor aggressiveness in renal cancer

Glucose metabolism is integral to cellular energy production, and its disruption is a key driver in the progression of renal cancer, particularly ccRCC. In ccRCC, HIFs orchestrate a metabolic reprogramming that boosts tumor aggressiveness. HIF-1 α not only promotes glucose uptake by stimulating GLUT transporter proteins, but it also suppresses mitochondrial respiration through the regulation of microRNAs such as miR-210 [33]. Conversely, HIF-2 α governs genes linked to glycolysis and interacts with pivotal oncogenes like MYC and P53, while upregulating cell cycle regulators [34]. This broad spectrum of HIF-2 α activity is central to ccRCC pathogenesis [35]. Moreover, the advent of selective HIF-2 α inhibitors has shown encouraging outcomes in ccRCC xenograft models, suggesting potential clinical utility [36]. HIF-1 α drives the preference for aerobic glycolysis which is a defining feature of ccRCC. It is evident from increased lactate production coupled with lower pyruvate flux into mitochondria, leading to suppressed TCA cycle activity and reduced ATP generation [37]. HIF-1 α amplifies these changes by inhibiting pyruvate dehydrogenase, which prevents pyruvate from converting into acetyl-CoA and drives lactate accumulation [38]. HIF-1 α also promote the expression of key glycolytic enzymes, including HK, neuron-specific enolase (NSE), PGK, and pyruvate kinase (PK), thereby reinforcing the Warburg effect in ccRCC [39, 40]. Among these enzymes, HK2 is of particular importance; its overexpression correlates with advanced tumor progression, lymph node metastasis, and poorer survival in renal cancer patients. HK2 also stands out as an independent risk factor for renal cell carcinoma, with studies linking its elevated expression to immune cell infiltration that impacts tumor progression and prognosis [41].

Glycolysis involves breaking glucose down into pyruvate [42]. Pyruvate enters the TCA cycle under aerobic conditions, supporting ATP generation along with the production of reduced

nicotinamide adenine dinucleotide (NADH) and reduced flavin adenine dinucleotide (FADH₂) [43]. Under anaerobic conditions, pyruvate is converted into lactate by fermentation, yielding ATP [44]. In ccRCC, another essential metabolic pathway is the pentose phosphate pathway, which generates glucose for lipid metabolism and nucleic acid synthesis by producing reduced nicotinamide adenine dinucleotide phosphate (NADPH) and ribose 5-phosphate [45]. The pentose phosphate pathway is significantly heightened in ccRCC, producing abundant NADPH necessary for preserving redox balance and safeguarding cancer cells from reactive oxygen species (ROS) damage [46]. This metabolic adjustment helps tumor cells mitigate oxidative stress and limit ROS-induced harm [47]. Additionally, the pentose phosphate pathway furnishes the five-carbon sugars required for nucleotide production, meeting the heightened demands of rapidly proliferating tumor cells [48]. Significant modifications also occur within the TCA cycle in ccRCC, contributing to its distinctive metabolic profile. Enzymes essential for refilling metabolic intermediates from other pathways are frequently reduced [49]. Of note, citrate and cis-aconitate are found in higher concentrations in ccRCC's TCA cycle, whereas

malate and fumarate are markedly decreased [15]. The drop in fumarate and malate is largely tied to the inhibition of succinate dehydrogenase (SDH), a mechanism that continuously diminishes fumarate and, by extension, malate [49]. This observation contradicts the typical notion of tumor tissues maintaining abundant fumarate. Overall, the abnormal regulation of glucose metabolism, primarily steered by HIF-1 α and HIF-2 α , plays a pivotal role in ccRCC aggressiveness. Collectively, overexpression of glycolytic enzymes, and alterations in the TCA cycle and pentose phosphate pathway grant ccRCC a metabolic advantage that propels its progression (Figure 2).

Warburg effect fosters metabolic supremacy-driven tumor aggressiveness in renal cancer

The Warburg effect describes cancer cells favoring glycolysis over mitochondrial respiration even with sufficient oxygen, resulting in limited ATP production and elevated lactate levels [12]. Recent research underscores the Warburg effect as a key driver of tumor growth in ccRCC. In this regard, ccRCC exhibits marked




Figure 1. Hypoxia signaling induces metabolic supremacy in renal cancer. In response to poor oxygen supply in growing tumors in ccRCC, hypoxia signaling, primarily HIF-2 α mediated one, gets activated leading to activation of plethora of genes involved in metabolic reprogramming. This results in onset of angiogenesis, tumor aggressiveness and progression, and metastatic spread of the disease.

metabolic reprogramming, particularly in the TCA cycle. Such changes include impaired pyruvate transport into mitochondria and dysfunction of enzymes like fumarate hydratase (FH) and SDH, leading to metabolic imbalances [50]. In addition, citrate and aconitate levels rise while succinate and malic acid levels fall [49], and the downregulation of enzymes that normally replenish the TCA cycle compounds these disruptions [50]. As a result, oxidative phosphorylation, the primary ATP-producing process, is severely compromised in ccRCC [49]. A hallmark of ccRCC is mutation of the von Hippel-Lindau (VHL) gene, which stabilizes HIF-1 α . This stabilization alters cellular metabolism by impeding glucose oxidation, reducing mitochondrial pyruvate uptake, and lowering the efficiency of the electron transport chain [51]. Oxidative phosphorylation impairment is further aggravated by the reduced expression of complex V and the downregulation of peroxisome proliferator-activated receptor gamma coactivator-1 α (PGC-1 α), a key regulator of mitochondrial biogenesis and respiration [52]. Decreased PGC-1 α delays mitochondrial respiration, lowers the expression of mitochondrial transcription factor A (TFAM), and correlates with worse clinical outcomes in ccRCC [27]. Besides HIF-1 α , HIF-2 α also influences oxidative phosphorylation activity in ccRCC by inducing antioxidant gene expression, thereby minimizing ROS accumulation, preventing DNA damage, and enhancing tumor cell survival [53]. These oxidative phosphorylation-related metabolic changes profoundly affect both tumor progression and treatment strategies, with selective HIF-2 α inhibitors showing notable therapeutic success in ccRCC [54]. In ccRCC, elevated glycolysis and declining oxygen levels in the tumor microenvironment raise the NADH/NAD⁺ ratio, disrupting redox balance and promoting ROS production [55]. Cancer cells counteract increased ROS by employing antioxidant defenses such as the thioredoxin and glutathione pathways [10]. Moreover, higher ROS levels stabilize nuclear factor erythroid 2-like 2 (NRF2), a critical antioxidant response regulator. This occurs when KEAP1, normally responsible for targeting NRF2 for proteasomal degradation, becomes oxidized and can no longer carry out its negative regulatory function [55]. Since excessive ROS can overwhelm a tumor cell's antioxidant defenses, these cells often bolster these defense mechanisms to ensure survival [56]. In ccRCC, HIF-2 α supports these processes by inducing antioxidant genes that further lower ROS levels, reduce DNA damage, and enhance tumor cell viability [53]. Overall, metabolic supremacy, particularly the interplay between disrupted oxidative phosphorylation and glycolysis, is key to tumor aggressiveness in renal cancer (**Figure 2**).

Dysregulated Amino acid metabolism promotes metabolic supremacy-driven tumor aggressiveness in renal cancer

Glutamine plays a key role in sustaining cellular redox homeostasis in normal cells by serving as a precursor for both α -ketoglutarate (α -KG) and glutathione, which are central to maintaining intracellular redox balance and facilitating amino acid synthesis [57]. It is transported into cells through specific transporter proteins such as Solute Carrier Family 1 Member 5 (SLC1A5) [58], after which the enzyme glutaminase (GLS) converts glutamine to glutamate [59]. Besides contributing to protein synthesis, glutamine underpins several metabolic and biosynthetic processes, including nucleotide and hexosamine production and asparagine formation, and is vital for managing oxidative stress and regulating other essential amino acids [60]. GLS activity is directly linked to both cell proliferation and cancer progression, making it an attractive target for anticancer strategies; indeed, inhibiting GLS expression or activity has been shown to restrict tumor growth [61]. In ccRCC, metabolic reprogramming heightens reliance on glutamine to power pathways that drive accelerated

tumor growth. Meeting the elevated metabolic demands of these aggressive cancer cells depends on glutamine [62]. Furthermore, an overabundance of glutamic acid disrupts cystine uptake, generating imbalances in ROS and impairing T cell functionality, thus promoting an immunosuppressive tumor microenvironment (**Figure 2**) [13]. A hallmark of ccRCC is the increased expression of SLC7A5, an amino acid transporter regulated by HIF-2 α , which enhances glutamine uptake [63]. Once inside the cell, glutamine is first converted to glutamate by GLS and then metabolized to α -KG through the action of glutamate dehydrogenase (GDH), providing crucial carbon for the TCA cycle and supporting cell survival [64]. Glutamate can also be channeled into reductive carboxylation to yield isocitrate, which is then used to produce acetyl coenzyme A for lipid biosynthesis [57]. The elevated glutamine content found in ccRCC correlates with increased glutamate production, a key mechanism for neutralizing ROS [65]. In addition, glutamine drives glycolysis, bolsters proliferation and immortalization, and hinders apoptosis by suppressing thioredoxin-interacting protein, further underscoring the therapeutic relevance of targeting glutamine metabolism in ccRCC.

Tryptophan also critically modulates T cell-mediated immune responses to tumors. However, its excessive oxidation through the kynurenine pathway triggers T cell dysfunction and allows tumor cells to evade immune detection [66]. Metabolites in the kynurenine pathway actively suppress T cell activation, exacerbating immune escape. Within tumor-draining lymph nodes, elevated indoleamine 2,3-dioxygenase (IDO) activity creates an immunosuppressive milieu by prompting dendritic cells to inhibit T cells, thereby interfering with antigen recognition and immune function [67]. In ccRCC, immune checkpoint dysregulation is associated with increased IDO expression, which depletes tryptophan and activates the kynurenine pathway, ultimately supporting tumor survival by countering interferon-alpha (IFN- α) therapy and fostering immune suppression (**Figure 2**) [68, 69]. In addition, IDO overexpression has been closely tied to cancer metastasis: research in lung cancer cells reveals that higher IDO levels improve cell viability, whereas IDO inhibition diminishes it. In mouse models, administering lung cancer cells overexpressing IDO leads to more frequent metastases in the brain, liver, and bone [70]. These observations pinpoint IDO as a promising therapeutic target in ccRCC and other malignancies, warranting further exploration. Arginine metabolism is similarly disrupted in ccRCC, involving abnormalities in arginine transporters and metabolic enzymes, including arginase and arginine succinate synthase 1 (ASS1). Tumor cells often display reduced or missing ASS1 expression, an enzyme needed to convert citrulline into arginine, forcing cancer cells to rely on external arginine. Proteomic analyses of ccRCC biopsies support this dependency. Targeting arginine metabolism thus offers a potential therapeutic approach, as depriving tumor cells of this vital nutrient can suppress cancer progression. Research has shown that eliminating arginine selectively induces cell toxicity in ASS1-deficient tumors [71]. In summary, disrupted amino acid metabolism contribute to metabolic supremacy and tumor aggressiveness in renal cancer (**Figure 2**).

Elevated lipid metabolism fuels metabolic supremacy-driven tumor aggressiveness in renal cancer

In ccRCC, disruptions in lipid metabolism frequently occur, markedly influencing the tumor's aggressive characteristics (**Figure 2**) [72]. Amplified lipid synthesis and storage, along with reduced lipid oxidation and utilization mark the metabolic supremacy, leading to accumulation of substantial levels of cholesterol, fatty acids, and triglycerides [73, 74], which support membrane formation and cell proliferation while limiting fatty




Figure 2. Metabolic supremacy drives tumor aggressiveness in renal cancer. Metabolic supremacy marked by elevated glucose, fatty acid, glutamine and tryptophan metabolism drives tumor aggressiveness in renal cancer. Glucose metabolism via glycolysis rather than oxidative phosphorylation (Warburg effect) fuels tumor proliferation and progression. Increased fatty acid and cholesterol uptake but reduced fatty acid oxidation leads to lipid accumulation which also serve as fuel for cancer progression. Meanwhile, enhanced glutamine metabolism counteracts oxidative stress via upregulating glutathione levels. Tumor cells limit arginine metabolism and rely on external arginine sources for rapid growth. Lastly, elevated tryptophan metabolism via IDO contribute to immune evasion.

acid β -oxidation. Driven by increased expression of lipoprotein receptors such as very low-density lipoprotein receptor (VLDL-R) and scavenger receptor B1 (SR-B1), enhanced cholesterol uptake is a major contributor to this imbalance [75, 76]. Moreover, HIF-2 α escalates lipid storage by activating hypoxia-induced lipid droplet-associated protein (HILPDA), leading to the selective enrichment of polyunsaturated lipids [77]. Overexpression of fatty acid synthase (FAS) represent another hallmark of ccRCC, leading to elevated fatty acid biosynthesis [39]. However, in contrast to other malignancies, ccRCC exhibits diminished fatty acid oxidation, primarily due to decreased levels of carnitine palmitoyltransferase 1A (CPT1A), an enzyme essential for fatty acid degradation [78]. In addition, ccRCC cells display higher amounts of fatty acid desaturase 1 (FADS1), which fosters polyunsaturated fatty acid (PUFA) production [72]. These PUFAs are crucial substrates for lipid peroxidation, a defining feature of ferroptosis. Research points to acyl-CoA synthetase long-chain family member 4 (ACSL4) as a key regulator of ferroptotic

sensitivity [79]. Once ACSL4 is active, lysophosphatidylcholine acyltransferase 3 (LPCAT3) facilitates ferroptotic signaling by integrating acyl groups into phospholipids like phosphatidylcholine and phosphatidylethanolamine. Nevertheless, ferroptosis can also arise through ACSL4-independent pathways, and although inducing ferroptosis is being investigated as a potential cancer therapy, its exact function in tumor biology remains a subject of ongoing exploration [80]. Lipid metabolism is closely interwoven with glucose metabolism. Enzymes such as glycerol kinase and α -phosphoglycerol dehydrogenase link glycerol and glucose pathways, yielding dihydroxyacetone phosphate [45]. In conditions where glucose is abundant, acetyl coenzyme A, derived from glucose metabolism, together with NADPH and protons from the pentose phosphate pathway, is converted into fatty acids via FAS. Metabolism of phospholipids and ketone bodies also contributes significantly to lipid metabolic processes [81]. Beyond their role as energy sources, lipids function as signaling molecules that regulate cell growth and proliferation [82]. Given its profound impact on

ccRCC progression, lipid metabolism is increasingly regarded as a promising target for therapeutic intervention.

Targeting metabolic supremacy in renal cancer

Renal cancer is often characterized by genetic disruptions that create a hypoxic environment, predominantly through mutations in the VHL gene [83]. These mutations lead to an accumulation of HIF- α , which in turn stimulates the production of vascular endothelial growth factors (VEGFs) [84]. Historically, therapies for ccRCC have focused on antiangiogenic measures, such as VEGF receptor inhibitors or agents directly targeting VEGF, including sunitinib [85]. However, these approaches have demonstrated limited clinical benefits and are frequently accompanied by adverse effects. Because ccRCC involves significant metabolic reprogramming, attention has turned toward enzymes and proteins that modulate these altered pathways, with the goal of selectively eradicating tumor cells while sparing normal tissue [86]. One key aspect of this shift is the heightened reliance of ccRCC on glycolysis. Limiting glycolysis has been proposed as a viable therapeutic intervention, in line with strategies employed in hepatocellular carcinoma [87]. Although early clinical findings indicate that restricting glycolysis could curb cancer progression, large-scale studies are still needed to verify its clinical value [88]. Delving deeper into the metabolic deviations characterizing ccRCC may thus lead to more effective and less toxic treatments for patients [89].

A particularly auspicious tactic targets the HIF-2 α signaling pathway, a major downstream effector of the frequently mutated VHL tumor suppressor gene in ccRCC [90]. HIF-2 α governs critical functions such as angiogenesis, cell proliferation, and metabolism, each substantially influencing tumor expansion and metastasis. Formerly deemed “undruggable” [91], HIF-2 α has recently been shown to possess a structural vulnerability in its PAS-B domain. This discovery spurred the development of initial HIF-2 α inhibitors like PT2399 and PT2385, which alter the PAS-B domain’s shape to block the formation of the HIF-2 α /HIF-1 β complex [92]. PT2399 has even surpassed sunitinib in certain models and remained effective against sunitinib-resistant tumors, although prolonged therapy can induce resistance via mutations in the binding site or HIF-1 β [93], and it does not fully suppress all HIF-2 α target genes. PT2385, meanwhile, showed a favorable safety profile and minimal toxicity in phase I trials, with a complete response observed in 2% of patients, partial responses in 12%, and stable disease in 52% [94]. In response to the shortcomings of these first-generation inhibitors, second-generation HIF-2 α antagonists have been created, such as PT2977 (also known as MK-6482 or belzutifan) [95]. PT2977 targets a region adjacent to the PAS-B domain, triggering a conformational change that disrupts gene interactions. It also boasts low lipophilicity, high oral bioavailability, and an encouraging safety profile. In a phase II trial, a daily dose of 120 mg resulted in a 49% objective response rate among ccRCC patients, with primarily mild and manageable side effects [96, 97]. Consequently, newer HIF-2 α inhibitors, particularly PT2977, have emerged as promising targeted therapeutics for ccRCC, offering improved potency and tolerability compared to older treatments.

Cancer cells also exploit glutamine metabolism to support energy generation, redox balance, and the synthesis of essential macromolecules. In ccRCC, GLS replenishes the TCA cycle and modestly drives cell proliferation [12]. The GLS inhibitor CB-839 has demonstrated strong anticancer activity in preclinical models. In animal studies, combining CB-839 with Everolimus, an mTOR inhibitor commonly used in ccRCC, enhanced antitumor efficacy. Clinical exploration of this combination, however, remains sparse, emphasizing the need for more extensive safety and effectiveness

data [98, 99]. Furthermore, some cancers exhibit elevated arginine dependence due to deficient ASS1, thereby increasing their reliance on external arginine [100]. Since arginine is pivotal for nitric oxide production and protein biosynthesis, reducing circulating arginine through ADI-PEG20 (a PEG-conjugated arginine deaminase) has been proposed as a strategy to constrain tumor growth in ccRCC. That said, the re-expression of ASS1 may diminish its impact [12]. Clinical data point to good tolerance of ADI-PEG20 and suggest it may overcome drug resistance in cancers that rely heavily on arginine [101]. Additionally, encouraging results have been reported for ADI-PEG20 in non-small cell lung cancer, acute myeloid leukemia, and uveal melanoma [102, 103]. Further research will be crucial for refining combination regimens and determining how best to avert resistance. The enzyme IDO degrades tryptophan via the kynurenine pathway, contributing to an immunosuppressive tumor environment by lowering tryptophan levels, thereby impeding T-cell function and promoting metastasis [69]. IDO inhibition has thus become an appealing immunotherapeutic avenue. The selective IDO inhibitor Epacadostat was found in preclinical studies to enhance the response of tumor-specific T cells [104], though clinical outcomes have been mixed, showing toxicity issues and moderate efficacy at best. Early-phase trials combining Epacadostat with the PD-1 inhibitor pembrolizumab showed limited but notable antitumor responses in advanced solid tumors; however, more research is needed to fully establish its clinical benefit [105]. Meanwhile, Navoximod has displayed acceptable tolerability and moderate bioavailability at 800 mg twice daily. Although Navoximod monotherapy showed limited impact, pairing it with Atezolizumab produced encouraging safety profiles and measurable antitumor activity, with ongoing trials investigating its broader therapeutic potential [106-108]. Additional IDO inhibitors, such as KHK2455, LY3381916, and MK-7162, are in clinical studies to evaluate their safety and efficacy [67]. Moving forward, it will be essential to refine combination therapies and dosing strategies for IDO-targeted treatments. When considered alongside approaches like glycolysis inhibition, HIF-2 α antagonism, and interventions involving lipid, glutamine, and arginine pathways, the growing range of metabolic therapies holds real promise for improving outcomes in ccRCC.

Genetic alterations in ccRCC also impact lipid metabolism, which is integral to tumor proliferation. FAS overexpression, a common finding, elevates intracellular fatty acid concentrations, fueling cancer growth and affecting post-translational modifications. Fatty acids are critical for both energy production and the maintenance of redox balance [109]. These insights have led researchers to propose inhibiting fatty acid synthesis as a therapeutic approach, supported by studies correlating greater FAS expression with higher tumor aggressiveness and worse clinical outcomes [110]. Preclinical evaluations indicate that the FAS inhibitor C75 can limit ccRCC cell proliferation and aggressiveness [111]. Another agent, TVB-2640, has shown promise in clinical contexts. A phase I investigation reported reduced fatty acid production in patients with non-small cell lung cancer, and follow-up trials in breast and ovarian cancers confirmed its efficacy and generally mild dermatological and ocular side effects [112]. TVB-2640 is currently undergoing evaluation in numerous cancer trials, including studies focused on ccRCC, suggesting that FAS inhibitors could eventually play a valuable role in treating this disease.

Conclusion and future prospect

Renal cancer is marked by a striking reconfiguration of its metabolic processes, fundamentally shifting how these tumor cells manage energy production and maintain redox balance.

Hypoxia-driven metabolic supremacy accompanied by metabolic shift from oxidative phosphorylation to glycolysis propels rapid tumor proliferation and aggressiveness. At the same time, the pentose phosphate pathway undergoes upregulation, boosting the generation of NADPH, which provides critical protection against ROS and helps safeguard nucleotides from potential damage. Excess lactate in the tumor environment further promotes an immunosuppressive milieu, thereby advancing tumor cell migration and invasion. ccRCC cells also exhibit significant shifts in lipid metabolism, with elevated lipid synthesis and utilization coinciding with the downregulation of lipid oxidation. ccRCC cells maintain high glutamine uptake despite such metabolic alterations, which underpins fatty acid production and serves as a buffer against oxidative stress by mitigating ROS. Taken together, these wide-ranging metabolic changes underscore a complex biochemical supremacy in ccRCC, emphasizing the necessity for continued research to clarify the nuances of these pathways and identify viable therapeutic targets. In view of the limited success rates of current treatments, tailored approaches that selectively disrupt vital metabolic pathways in cancer cells by leveraging the unique metabolic characteristics of ccRCC, open new possibilities for drug development. The potential benefits of intervening in tumor metabolism likely surpass the drawbacks and challenges associated with conventional therapies such as, potential off-target effects on other rapidly dividing cells and variability arising from distinct mutation profiles. For instance, metabolic inhibitors could ultimately present a safer option than standard anti-angiogenic therapies, which are frequently linked to cardiovascular side effects. Moreover, advances in metabolomics continue to reveal novel metabolic targets, paving the way for precision therapies specifically suited to ccRCC. As research progresses, it is anticipated that new treatments aimed at metabolic supremacy will broaden the range of viable options for patients with ccRCC, offering both improved outcomes and a more personalized approach to care.

Acknowledgements

None.

Ethical policy

Non applicable.

Availability of data and materials

All data generated or analysed during this study are included in this publication.

Author contributions

Reham Gholam contributed to design of the work, data collection, and drafting the article. Muhammad Khalilzad was devoted to critical revision and final submission of the article.

Competing interests

The authors declare no competing interests.

Funding

None.

References

1. Aden D, Sureka N, Zaheer S, Chaurasia JK, Zaheer S: Metabolic

- Reprogramming in Cancer: Implications for Immunosuppressive Microenvironment. *Immunology* 2025, 174(1): 30-72.
2. Nong S, Han X, Xiang Y, Qian Y, Wei Y, Zhang T, Tian K, Shen K, Yang J, Ma X: Metabolic reprogramming in cancer: Mechanisms and therapeutics. *MedComm (2020) 2023*, 4(2): e218.
 3. Peixoto C, Martins M: Kidney Cancer Biomarker Selection Using Regularized Survival Models. *Cells* 2022, 11(15): 2311.
 4. Siddiqi A, Rani M, Bansal P, Rizvi MMA: Renal cell carcinoma management: A step to nano-chemoprevention. *Life Sci* 2022, 308: 120922.
 5. Hu SL, Chang A, Perazella MA, Okusa MD, Jaimes EA, Weiss RH: The Nephrologist's Tumor: Basic Biology and Management of Renal Cell Carcinoma. *J Am Soc Nephrol* 2016, 27(8): 2227-2237.
 6. Szymanski Ł, Helbrecht I, Fiedorowicz M, Matak D, Bartnik E, Golik P, Szczylik C, Czarnecka AM: [Cancer stem cells in renal carcinoma]. *Postepy Biochem* 2019, 65(2):95-102.
 7. Barata PC, Rini BI: Treatment of renal cell carcinoma: Current status and future directions. *CA Cancer J Clin* 2017, 67(6): 507-524.
 8. Motzer RJ, Powles T, Burotto M, Escudier B, Bourlon MT, Shah AY, Suárez C, Hamzaj A, Porta C, Hocking CM et al: Nivolumab plus cabozantinib versus sunitinib in first-line treatment for advanced renal cell carcinoma (CheckMate 9ER): long-term follow-up results from an open-label, randomised, phase 3 trial. *Lancet Oncol* 2022, 23(7): 888-898.
 9. Rini BI, Plimack ER, Stus V, Gafanov R, Hawkins R, Nosov D, Pouliot F, Alekseev B, Soulières D, Melichar B et al: Pembrolizumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma. *N Engl J Med* 2019, 380(12): 1116-1127.
 10. Faubert B, Solmonson A: Metabolic reprogramming and cancer progression. *Science* 2020, 368(6487): eaaw5473.
 11. Warburg O: On the origin of cancer cells. *Science* 1956, 123(3191): 309-314.
 12. Wettersten HI, Aboud OA, Lara PN Jr, Weiss RH: Metabolic reprogramming in clear cell renal cell carcinoma. *Nat Rev Nephrol* 2017, 13(7): 410-419.
 13. Cassim S, Pouyssegur J: Tumor Microenvironment: A Metabolic Player that Shapes the Immune Response. *Int J Mol Sci* 2019, 21(1): 157.
 14. Jonasch E, Walker CL, Rathmell WK: Clear cell renal cell carcinoma ontogeny and mechanisms of lethality. *Nat Rev Nephrol* 2021, 17(4): 245-261.
 15. Hakimi AA, Reznik E, Lee CH, Creighton CJ, Brannon AR, Luna A, Aksoy BA, Liu EM, Shen R, Lee W et al: An Integrated Metabolic Atlas of Clear Cell Renal Cell Carcinoma. *Cancer Cell* 2016, 29(1): 104-116.
 16. Fu Q, Xu L, Wang Y, Jiang Q, Liu Z, Zhang J, Zhou Q, Zeng H, Tong S, Wang T et al: Tumor-associated Macrophage-derived Interleukin-23 Interlinks Kidney Cancer Glutamine Addiction with Immune Evasion. *Eur Urol* 2019, 75(5): 752-763.
 17. Linehan WM, Schmidt LS, Crooks DR, Wei D, Srinivasan R, Lang M: The Metabolic Basis of Kidney Cancer. *Cancer Discov* 2019, 9(8): 1006-1021.
 18. Chitrakar A, Budda SA, Henderson JG, Axtell RC, Zenewicz LA: E3 Ubiquitin Ligase Von Hippel-Lindau Protein Promotes Th17 Differentiation. *J Immunol* 2020, 205(4): 1009-1023.
 19. Chappell JC, Payne LB, Rathmell WK: Hypoxia, angiogenesis, and metabolism in the hereditary kidney cancers. *J Clin Invest* 2019, 129(2): 442-451.
 20. Lee P, Chandel NS: Cellular adaptation to hypoxia through hypoxia inducible factors and beyond. *Nat Rev Mol Cell Biol* 2020, 21(5): 268-283.
 21. Al Tameemi W, Dale TP, Al-Jumaily RMK, Forsyth NR: Hypoxia-Modified Cancer Cell Metabolism. *Front Cell Dev Biol* 2019, 7: 4.
 22. Semenza GL: HIF-1 mediates metabolic responses to intratumoral hypoxia and oncogenic mutations. *J Clin Invest* 2013, 123(9):

- 3664-3671.
23. Choudhry H, Harris AL: Advances in Hypoxia-Inducible Factor Biology. *Cell Metab* 2018, 27(2): 281-298.
 24. Jaakkola P, Mole DR, Tian YM, Wilson MI, Gielbert J, Gaskell SJ, von Kriegsheim A, Hebestreit HF, Mukherji M, Schofield CJ et al: Targeting of HIF- α to the von Hippel-Lindau ubiquitylation complex by O₂-regulated prolyl hydroxylation. *Science* 2001, 292(5516): 468-472.
 25. Motzer RJ, Banchereau R, Hamidi H, Powles T, McDermott D, Atkins MB, Escudier B, Liu LF, Leng N, Abbas AR et al: Molecular Subsets in Renal Cancer Determine Outcome to Checkpoint and Angiogenesis Blockade. *Cancer Cell* 2020, 38(6): 803-817.e804.
 26. Schönenberger D, Harlander S, Rajski M, Jacobs RA, Lundby AK, Adlesic M, Hejhal T, Wild PJ, Lundby C, Frew IJ: Formation of Renal Cysts and Tumors in Vhl/Trp53-Deficient Mice Requires HIF1 α and HIF2 α . *Cancer Res* 2016, 76(7): 2025-2036.
 27. LaGory EL, Wu C, Taniguchi CM, Ding CC, Chi JT, von Eyben R, Scott DA, Richardson AD, Giaccia AJ: Suppression of PGC-1 α Is Critical for Reprogramming Oxidative Metabolism in Renal Cell Carcinoma. *Cell Rep* 2015, 12(1): 116-127.
 28. Dibble CC, Cantley LC: Regulation of mTORC1 by PI3K signaling. *Trends Cell Biol* 2015, 25(9): 545-555.
 29. Wee S, Wiederschain D, Maira SM, Loo A, Miller C, deBeaumont R, Stegmeier F, Yao YM, Lengauer C: PTEN-deficient cancers depend on PIK3CB. *Proc Natl Acad Sci U S A* 2008, 105(35): 13057-13062.
 30. Yang P, Cornejo KM, Sadow PM, Cheng L, Wang M, Xiao Y, Jiang Z, Oliva E, Jozwiak S, Nussbaum RL et al: Renal cell carcinoma in tuberous sclerosis complex. *Am J Surg Pathol* 2014, 38(7): 895-909.
 31. Yu HB, Kielczewska A, Rozek A, Takenaka S, Li Y, Thorson L, Hancock REW, Guarna MM, North JR, Foster LJ et al: Sequestosome-1/p62 is the key intracellular target of innate defense regulator peptide. *J Biol Chem* 2009, 284(52): 36007-36011.
 32. Düvel K, Yecies JL, Menon S, Raman P, Lipovsky AI, Souza AL, Triantafellow E, Ma Q, Gorski R, Cleaver S et al: Activation of a metabolic gene regulatory network downstream of mTOR complex 1. *Mol Cell* 2010, 39(2): 171-183.
 33. Ivan M, Huang X: miR-210: fine-tuning the hypoxic response. *Adv Exp Med Biol* 2014, 772: 205-227.
 34. Schödel J, Grampp S, Maher ER, Moch H, Ratcliffe PJ, Russo P, Mole DR: Hypoxia, Hypoxia-inducible Transcription Factors, and Renal Cancer. *Eur Urol* 2016, 69(4): 646-657.
 35. Kondo K, Kim WY, Lechpammer M, Kaelin WG Jr: Inhibition of HIF2 α is sufficient to suppress pVHL-defective tumor growth. *PLoS Biol* 2003, 1(3): E83.
 36. Courtney KD, Ma Y, Diaz de Leon A, Christie A, Xie Z: HIF-2 Complex Dissociation, Target Inhibition, and Acquired Resistance with PT2385, a First-in-Class HIF-2 Inhibitor, in Patients with Clear Cell Renal Cell Carcinoma. *Clin Cancer Res* 2020, 26(4): 793-803.
 37. Kim JW, Tchernyshyov I, Semenza GL, Dang CV: HIF-1-mediated expression of pyruvate dehydrogenase kinase: a metabolic switch required for cellular adaptation to hypoxia. *Cell Metab* 2006, 3(3): 177-185.
 38. Infantino V, Santarsiero A: Cancer Cell Metabolism in Hypoxia: Role of HIF-1 as Key Regulator and Therapeutic Target. *Int J Mol Sci* 2021, 22(11): 5703.
 39. Xie H, Simon MC: Oxygen availability and metabolic reprogramming in cancer. *J Biol Chem* 2017, 292(41): 16825-16832.
 40. Nakazawa MS, Keith B, Simon MC: Oxygen availability and metabolic adaptations. *Nat Rev Cancer* 2016, 16(10): 663-673.
 41. Liu C, Li H, Huang H, Zheng P, Li Z: The Correlation of HK2 Gene Expression with the Occurrence, Immune Cell Infiltration, and Prognosis of Renal Cell Carcinoma. *Dis Markers* 2022, 2022: 1452861.
 42. Adeva-Andany MM, Pérez-Felpete N, Fernández-Fernández C, Donapetry-García C, Pazos-García C: Liver glucose metabolism in humans. *Biosci Rep* 2016, 36(6): e00416.
 43. Rui L: Energy metabolism in the liver. *Compr Physiol* 2014, 4(1): 177-197.
 44. Han HS, Kang G, Kim JS, Choi BH, Koo SH: Regulation of glucose metabolism from a liver-centric perspective. *Exp Mol Med* 2016, 48(3): e218.
 45. Wanders RJ, Vreken P, Ferdinandusse S, Jansen GA, Waterham HR, van Roermund CW, Van Grunsven EG: Peroxisomal fatty acid alpha- and beta-oxidation in humans: enzymology, peroxisomal metabolite transporters and peroxisomal diseases. *Biochem Soc Trans* 2001, 29(Pt 2): 250-267.
 46. Nogueira V, Hay N: Molecular pathways: reactive oxygen species homeostasis in cancer cells and implications for cancer therapy. *Clin Cancer Res* 2013, 19(16): 4309-4314.
 47. Aykin-Burns N, Ahmad IM, Zhu Y, Oberley LW, Spitz DR: Increased levels of superoxide and H₂O₂ mediate the differential susceptibility of cancer cells versus normal cells to glucose deprivation. *Biochem J* 2009, 418(1): 29-37.
 48. Jiang P, Du W, Wu M: Regulation of the pentose phosphate pathway in cancer. *Protein Cell* 2014, 5(8): 592-602.
 49. Chakraborty S, Balan M, Sabarwal A, Choueiri TK, Pal S: Metabolic reprogramming in renal cancer: Events of a metabolic disease. *Biochim Biophys Acta Rev Cancer* 2021, 1876(1): 188559.
 50. Massari F, Ciccarese C, Santoni M, Brunelli M, Piva F, Modena A, Bimbatti D, Fantinel E, Santini D, Cheng L et al: Metabolic alterations in renal cell carcinoma. *Cancer Treat Rev* 2015, 41(9): 767-776.
 51. Wettersten HI, Hakimi AA, Morin D, Bianchi C, Johnstone ME, Donohoe DR, Trott JF, Aboud OA, Stirdivant S, Neri B et al: Grade-Dependent Metabolic Reprogramming in Kidney Cancer Revealed by Combined Proteomics and Metabolomics Analysis. *Cancer Res* 2015, 75(12): 2541-2552.
 52. Simonnet H, Alazard N, Pfeiffer K, Gallou C, Bérout C, Demont J, Bouvier R, Schägger H, Godinot C: Low mitochondrial respiratory chain content correlates with tumor aggressiveness in renal cell carcinoma. *Carcinogenesis* 2002, 23(5): 759-768.
 53. Yu Y, Yu Q, Zhang X: Allosteric inhibition of HIF-2 α as a novel therapy for clear cell renal cell carcinoma. *Drug Discov Today* 2019, 24(12): 2332-2340.
 54. Molina JR, Sun Y, Protopopova M, Gera S, Bandi M, Bristow C, McAfoos T, Morlacchi P, Ackroyd J: An inhibitor of oxidative phosphorylation exploits cancer vulnerability. *Nat Med* 2018, 24(7): 1036-1046.
 55. Zhu J, Thompson CB: Metabolic regulation of cell growth and proliferation. *Nat Rev Mol Cell Biol* 2019, 20(7): 436-450.
 56. Piskounova E, Agathocleous M, Murphy MM, Hu Z, Huddleston SE, Zhao Z, Leitch AM, Johnson TM, DeBerardinis RJ, Morrison SJ: Oxidative stress inhibits distant metastasis by human melanoma cells. *Nature* 2015, 527(7577): 186-191.
 57. Hensley CT, Wasti AT, DeBerardinis RJ: Glutamine and cancer: cell biology, physiology, and clinical opportunities. *J Clin Invest* 2013, 123(9): 3678-3684.
 58. Yoo HC, Park SJ, Nam M, Kang J, Kim K, Yeo JH, Kim JK, Heo Y, Lee HS, Lee MY et al: A Variant of SLC1A5 Is a Mitochondrial Glutamine Transporter for Metabolic Reprogramming in Cancer Cells. *Cell Metab* 2020, 31(2): 267-283.e212.
 59. Edwards DN, Ngwa VM, Raybuck AL, Wang S, Hwang Y, Kim LC, Cho SH, Paik Y, Wang Q, Zhang S et al: Selective glutamine metabolism inhibition in tumor cells improves antitumor T lymphocyte activity in triple-negative breast cancer. *J Clin Invest* 2021, 131(4): e140100.
 60. Pavlova NN, Zhu J, Thompson CB: The hallmarks of cancer

- metabolism: Still emerging. *Cell Metab* 2022, 34(3): 355-377.
61. Lobo C, Ruiz-Bellido MA, Aledo JC, Márquez J, Núñez De Castro I, Alonso FJ: Inhibition of glutaminase expression by antisense mRNA decreases growth and tumorigenicity of tumour cells. *Biochem J* 2000, 348 Pt 2(2): 257-261.
 62. Wise DR, Thompson CB: Glutamine addiction: a new therapeutic target in cancer. *Trends Biochem Sci* 2010, 35(8): 427-433.
 63. Onishi Y, Hiraiwa M, Kamada H, Iezaki T, Yamada T, Kaneda K, Hinoi E: Hypoxia affects Slc7a5 expression through HIF-2 α in differentiated neuronal cells. *FEBS Open Bio* 2019, 9(2): 241-247.
 64. Yang C, Sudderth J, Dang T, Bachoo RM, McDonald JG, DeBerardinis RJ: Glioblastoma cells require glutamate dehydrogenase to survive impairments of glucose metabolism or Akt signaling. *Cancer Res* 2009, 69(20): 7986-7993.
 65. Wettersten HI: Reprogramming of Metabolism in Kidney Cancer. *Semin Nephrol* 2020, 40(1): 2-13.
 66. Fiore A, Murray PJ: Tryptophan and indole metabolism in immune regulation. *Curr Opin Immunol* 2021, 70: 7-14.
 67. Platten M, Nollen EAA, Röhrig UF: Tryptophan metabolism as a common therapeutic target in cancer, neurodegeneration and beyond. *Nat Rev Drug Discov* 2019, 18(5): 379-401.
 68. Riesenberger R, Weiler C, Spring O, Eder M, Buchner A, Popp T, Castro M, Kammerer R, Takikawa O, Hatz RA et al: Expression of indoleamine 2,3-dioxygenase in tumor endothelial cells correlates with long-term survival of patients with renal cell carcinoma. *Clin Cancer Res* 2007, 13(23): 6993-7002.
 69. Trott JF, Kim J, Abu Aboud O, Wettersten H, Stewart B, Berryhill G, Uzal F, Hovey RC, Chen CH, Anderson K et al: Inhibiting tryptophan metabolism enhances interferon therapy in kidney cancer. *Oncotarget* 2016, 7(41): 66540-66557.
 70. Tang D, Yue L, Yao R, Zhou L, Yang Y, Lu L, Gao W: P53 prevent tumor invasion and metastasis by down-regulating IDO in lung cancer. *Oncotarget* 2017, 8(33): 54548-54557.
 71. Yao S, Janku F, Subbiah V: Phase I trial of ADI-PEG20 plus cisplatin in patients with pretreated metastatic melanoma or other advanced solid malignancies. *Br J Cancer* 2021, 124(9): 1533-1539.
 72. Heravi G, Yazdanpanah O, Podgorski I, Matherly LH, Liu W: Lipid metabolism reprogramming in renal cell carcinoma. *Cancer Metastasis Rev* 2022, 41(1): 17-31.
 73. Ganti S, Taylor SL, Abu Aboud O, Yang J, Evans C, Osier MV, Alexander DC, Kim K, Weiss RH: Kidney tumor biomarkers revealed by simultaneous multiple matrix metabolomics analysis. *Cancer Res* 2012, 72(14): 3471-3479.
 74. Gebhard RL, Clayman RV, Prigge WF, Figenshau R, Staley NA, Reese C, Bear A: Abnormal cholesterol metabolism in renal clear cell carcinoma. *J Lipid Res* 1987, 28(10): 1177-1184.
 75. Saito K, Arai E, Maekawa K, Ishikawa M, Fujimoto H, Taguchi R, Matsumoto K, Kanai Y, Saito Y: Lipidomic Signatures and Associated Transcriptomic Profiles of Clear Cell Renal Cell Carcinoma. *Sci Rep* 2016, 6: 28932.
 76. Wu G, Wang Q, Xu Y, Li J, Zhang H, Qi G, Xia Q: Targeting the transcription factor receptor LXR to treat clear cell renal cell carcinoma: agonist or inverse agonist? *Cell Death Dis* 2019, 10(6): 416.
 77. Zou Y, Palte MJ, Deik AA, Li H, Eaton JK: A GPX4-dependent cancer cell state underlies the clear-cell morphology and confers sensitivity to ferroptosis. *Nat Commun* 2019, 10(1): 1617.
 78. Du W, Zhang L, Brett-Morris A, Aguila B, Kerner J, Hoppel CL, Puchowicz M, Serra D, Herrero L: HIF drives lipid deposition and cancer in ccRCC via repression of fatty acid metabolism. *Nat Commun* 2017, 8(1): 1769.
 79. Kagan VE, Mao G, Qu F, Angeli JP, Doll S, Croix CS, Dar HH, Liu B, Tyurin VA, Ritov VB et al: Oxidized arachidonic and adrenic PEs navigate cells to ferroptosis. *Nat Chem Biol* 2017, 13(1): 81-90.
 80. Chen X, Li J, Kang R, Klionsky DJ: Ferroptosis: machinery and regulation. *Autophagy* 2021, 17(9): 2054-2081.
 81. Huang C, Freter C: Lipid metabolism, apoptosis and cancer therapy. *Int J Mol Sci* 2015, 16(1): 924-949.
 82. Hanahan D, Weinberg RA: Hallmarks of cancer: the next generation. *Cell* 2011, 144(5): 646-674.
 83. Hsieh JJ, Le VH, Oyama T, Ricketts CJ, Ho TH, Cheng EH: Chromosome 3p Loss-Orchestrated VHL, HIF, and Epigenetic Deregulation in Clear Cell Renal Cell Carcinoma. *J Clin Oncol* 2018, 36(36): Jco2018792549.
 84. Keith B, Johnson RS, Simon MC: HIF1 α and HIF2 α : sibling rivalry in hypoxic tumour growth and progression. *Nat Rev Cancer* 2011, 12(1): 9-22.
 85. Roskoski R Jr.: Vascular endothelial growth factor (VEGF) and VEGF receptor inhibitors in the treatment of renal cell carcinomas. *Pharmacol Res* 2017, 120: 116-132.
 86. Keefe D, Bowen J, Gibson R, Tan T, Okera M, Stringer A: Noncardiac vascular toxicities of vascular endothelial growth factor inhibitors in advanced cancer: a review. *Oncologist* 2011, 16(4): 432-444.
 87. Cassim S, Raymond VA, Dehbidi-Assadzadeh L, Lapierre P, Bilodeau M: Metabolic reprogramming enables hepatocarcinoma cells to efficiently adapt and survive to a nutrient-restricted microenvironment. *Cell Cycle* 2018, 17(7): 903-916.
 88. Abdel-Wahab AF, Mahmoud W, Al-Harizy RM: Targeting glucose metabolism to suppress cancer progression: prospective of anti-glycolytic cancer therapy. *Pharmacol Res* 2019, 150: 104511.
 89. DeBerardinis RJ, Keshari KR: Metabolic analysis as a driver for discovery, diagnosis, and therapy. *Cell* 2022, 185(15): 2678-2689.
 90. Kaelin WG Jr, Ratcliffe PJ: Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Mol Cell* 2008, 30(4): 393-402.
 91. Koehler AN: A complex task? Direct modulation of transcription factors with small molecules. *Curr Opin Chem Biol* 2010, 14(3): 331-340.
 92. Scheuermann TH, Li Q, Ma HW, Key J, Zhang L, Chen R, Garcia JA, Naidoo J, Longgood J, Frantz DE et al: Allosteric inhibition of hypoxia inducible factor-2 with small molecules. *Nat Chem Biol* 2013, 9(4): 271-276.
 93. Chen W, Hill H, Christie A, Kim MS, Holloman E, Pavia-Jimenez A, Homayoun F, Ma Y, Patel N, Yell P et al: Targeting renal cell carcinoma with a HIF-2 antagonist. *Nature* 2016, 539(7627): 112-117.
 94. Courtney KD, Infante JR, Lam ET, Figlin RA, Rini BI, Brugarolas J, Zojwalla NJ, Lowe AM, Wang K, Wallace EM et al: Phase I Dose-Escalation Trial of PT2385, a First-in-Class Hypoxia-Inducible Factor-2 α Antagonist in Patients With Previously Treated Advanced Clear Cell Renal Cell Carcinoma. *J Clin Oncol* 2018, 36(9): 867-874.
 95. Xu R, Wang K, Rizzi JP, Huang H, Grina JA, Schlachter ST, Wang B, Wehn PM, Yang H, Dixon DD et al: 3-[[1S,2S,3R)-2,3-Difluoro-1-hydroxy-7-methylsulfonylindan-4-yl]oxy-5-fluorobenzonitrile (PT2977), a Hypoxia-Inducible Factor 2 α (HIF-2 α) Inhibitor for the Treatment of Clear Cell Renal Cell Carcinoma. *J Med Chem* 2019, 62(15): 6876-6893.
 96. Jonasch E, Donskov F, Iliopoulos O, Rathmell WK, Narayan VK, Maughan BL, Oudard S, Else T, Maranchie JK, Welsh SJ et al: Belzutifan for Renal Cell Carcinoma in von Hippel-Lindau Disease. *N Engl J Med* 2021, 385(22): 2036-2046.
 97. Choueiri TK, Bauer TM, Papadopoulos KP, Plimack ER: Inhibition of hypoxia-inducible factor-2 α in renal cell carcinoma with belzutifan: a phase I trial and biomarker analysis. *Nat Med* 2021, 27(5): 802-805.
 98. Emberley E, Pan A, Chen J, Dang R, Gross M, Huang T, Li W, MacKinnon A, Singh D, Sotirovska N et al: The glutaminase inhibitor telaglenastat enhances the antitumor activity of signal

transduction inhibitors everolimus and cabozantinib in models of renal cell carcinoma. *PLoS One* 2021, 16(11): e0259241.

99. Delage B, Fennell DA, Nicholson L, McNeish I, Lemoine NR, Crook T, Szlosarek PW: Arginine deprivation and argininosuccinate synthetase expression in the treatment of cancer. *Int J Cancer* 2010, 126(12): 2762-2772.
100. Haines RJ, Pendleton LC, Eichler DC: Argininosuccinate synthase: at the center of arginine metabolism. *Int J Biochem Mol Biol* 2011, 2(1): 8-23.
101. Hall PE, Lewis R, Syed N, Shaffer R, Evanson J, Ellis S, Williams M: A Phase I Study of Pegylated Arginine Deiminase (Pegargiminase), Cisplatin, and Pemetrexed in Argininosuccinate Synthetase 1-Deficient Recurrent High-grade Glioma. *Clin Cancer Res* 2019, 25(9): 2708-2716.
102. Tomlinson BK, Thomson JA, Bomalaski JS, Diaz M, Akande T, Mahaffey N, Li T, Dutia MP, Kelly K, Gong IY et al: Phase I Trial of Arginine Deprivation Therapy with ADI-PEG 20 Plus Docetaxel in Patients with Advanced Malignant Solid Tumors. *Clin Cancer Res* 2015, 21(11): 2480-2486.
103. Abou-Alfa GK, Qin S, Ryoo BY, Lu SN, Yen CJ, Feng YH, Lim HY, Izzo F, Colombo M, Sarker D et al: Phase III randomized study of second line ADI-PEG 20 plus best supportive care versus placebo plus best supportive care in patients with advanced hepatocellular carcinoma. *Ann Oncol* 2018, 29(6): 1402-1408.
104. Jochems C, Fantini M, Fernando RI, Kwilas AR, Donahue RN, Lepone LM, Grenga I, Kim YS, Brechbiel MW, Gulley JL et al: The IDO1 selective inhibitor epacadostat enhances dendritic cell immunogenicity and lytic ability of tumor antigen-specific T cells. *Oncotarget* 2016, 7(25): 37762-37772.
105. Naing A, Powderly JD: Exploring the safety, effect on the tumor microenvironment, and efficacy of itacitinib in combination with epacadostat or pascalisib in advanced solid tumors: a phase I study. *J Immunother Cancer* 2022, 10(3): e004223.
106. Nayak-Kapoor A, Hao Z, Sadek R, Dobbins R, Marshall L, Vahanian NN, Jay Ramsey W, Kennedy E, Mautino MR, Link CJ et al: Phase Ia study of the indoleamine 2,3-dioxygenase 1 (IDO1) inhibitor navoximod (GDC-0919) in patients with recurrent advanced solid tumors. *J Immunother Cancer* 2018, 6(1): 61.
107. Ma S, Suchomel J, Yanez E, Yost E, Liang X, Zhu R, Le H, Siebers N, Joas L, Morley R et al: Investigation of the absolute bioavailability and human mass balance of navoximod, a novel IDO1 inhibitor. *Br J Clin Pharmacol* 2019, 85(8): 1751-1760.
108. Jung KH, LoRusso P, Burris H, Gordon M, Bang YJ, Hellmann MD: Phase I Study of the Indoleamine 2,3-Dioxygenase 1 (IDO1) Inhibitor Navoximod (GDC-0919) Administered with PD-L1 Inhibitor (Atezolizumab) in Advanced Solid Tumors. *Clin Cancer Res* 2019, 25(11): 3220-3228.
109. Angeles TS, Hudkins RL: Recent advances in targeting the fatty acid biosynthetic pathway using fatty acid synthase inhibitors. *Expert Opin Drug Discov* 2016, 11(12): 1187-1199.
110. Horiguchi A, Asano T, Asano T, Ito K, Sumitomo M, Hayakawa M: Fatty acid synthase over expression is an indicator of tumor aggressiveness and poor prognosis in renal cell carcinoma. *J Urol* 2008, 180(3): 1137-1140.
111. Horiguchi A, Asano T, Asano T, Ito K, Sumitomo M, Hayakawa M: Pharmacological inhibitor of fatty acid synthase suppresses growth and invasiveness of renal cancer cells. *J Urol* 2008, 180(2): 729-736.
112. Wang J, Lin W, Li R, Cheng H, Sun S, Shao F, Yang Y, Zhang L, Feng X, Gao S et al: The Deubiquitinase USP13 Maintains Cancer Cell Stemness by Promoting FASN Stability in Small Cell Lung Cancer. *Front Oncol* 2022, 12: 899987.



Copyright © 2025 Ann Urol Oncol. This work is licensed under a Creative Commons Attribution-NonCommercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) License.