

Metabolic Reprogramming in Kidney Cancer: Implications for Therapy

Elena Tena Edo¹

Cite this article: Edo TE: Metabolic Reprogramming in Kidney Cancer: Implications for Therapy. Ann Urol Oncol 2025, 8(2): 76-87. <https://doi.org/10.32948/auo.2025.06.02>

Abstract

Kidney cancer, particularly clear cell renal cell carcinoma (ccRCC), has emerged as a paradigm for cancer metabolic reprogramming, exhibiting distinctive alterations that drive tumor progression and therapeutic resistance. This comprehensive review synthesizes current knowledge of the molecular mechanisms underlying metabolic dysregulation in kidney cancer, with emphasis on the central role of VHL/HIF pathway activation and its downstream consequences on glycolysis, lipid metabolism, and glutamine utilization. We systematically analyze how pseudohypoxia-driven metabolic rewiring not only supports tumor bioenergetics and biosynthesis but also shapes an immunosuppressive microenvironment through metabolite-mediated crosstalk with stromal and immune cells. The review highlights groundbreaking therapeutic advances, including FDA-approved HIF-2 α inhibitors and emerging agents targeting glycolytic enzymes, glutaminase, and lipid metabolism, while addressing the challenges of metabolic plasticity and acquired resistance. Special attention is given to innovative combination strategies that pair metabolic modulators with immunotherapy or tyrosine kinase inhibitors, supported by preclinical rationale and clinical trial data. We further discuss cutting-edge technologies transforming the field - from hyperpolarized MRI for real-time metabolic imaging to AI-driven analysis of multi-omics datasets for patient stratification. By integrating fundamental science with translational applications, this review provides a framework for understanding kidney cancer as a metabolic disease and outlines future directions for targeted therapies, biomarker development, and personalized treatment approaches. The synthesis of these insights offers both a conceptual foundation and practical guidance for researchers and clinicians working to exploit metabolic vulnerabilities in kidney cancer.

Key words kidney cancer, ccRCC, VHL, metabolic reprogramming, therapeutics

1. Faculty of Health Sciences, International University of La Rioja, Av. de la Paz, 137, 26006 Logrono, La Rioja, Spain.

Correspondence: Elena Tena Edo (Faculty of Health Sciences, International University of La Rioja, Av. de la Paz, 137, 26006 Logrono, La Rioja, Spain; Email: elenatenaedo@gmail.com).

Introduction

Kidney cancer represents a compelling model for investigating the fundamental principles of cancer metabolism, with ccRCC exhibiting some of the most profound metabolic alterations observed in human malignancies [1]. The unique metabolic phenotype of kidney cancer stems from its distinctive genetic landscape, where inactivation of the Von Hippel-Lindau (VHL) tumor suppressor gene occurs in approximately 90% of sporadic ccRCC cases [2]. This genetic alteration triggers a cascade of molecular events that fundamentally reshape cellular metabolism, creating dependencies that differ markedly from normal renal epithelium [3]. The resulting metabolic reprogramming not only supports tumor growth and proliferation but also influences disease progression, treatment resistance, and interactions with the tumor microenvironment (TME) [4, 5]. This review provides a comprehensive examination of kidney cancer metabolism, exploring its molecular foundations, pathophysiological consequences, and emerging therapeutic opportunities.

At the heart of kidney cancer's metabolic transformation lies the constitutive activation of hypoxia-inducible factors (HIFs), particularly HIF-2 α , due to VHL loss [6]. Under normal oxygen conditions, VHL targets HIF- α subunits for proteasomal degradation, but in ccRCC, this regulatory mechanism fails, creating a state of pseudohypoxia regardless of actual oxygen availability [7]. HIF stabilization orchestrates a broad transcriptional program that upregulates glucose transporters (GLUT1, GLUT3) and glycolytic enzymes (HK2, PKM2, LDHA), while simultaneously suppressing mitochondrial oxidative phosphorylation [8] [9]. This metabolic shift, reminiscent of the Warburg effect but with unique kidney cancer-specific features, provides rapidly dividing tumor cells with essential biosynthetic precursors while maintaining redox homeostasis [10]. However, recent research has revealed that kidney cancer metabolism extends far beyond glycolysis, encompassing profound alterations in lipid, amino acid, and nucleotide metabolism that collectively sustain tumor growth and survival.

The lipid-rich phenotype of ccRCC represents one of its most distinctive metabolic features, visible histologically as cytoplasmic lipid droplets [11]. This characteristic results from coordinated increases in fatty acid uptake (mediated by CD36 and other transporters), enhanced de novo lipogenesis (through upregulation of FASN and ACC), and impaired lipid oxidation due to mitochondrial dysfunction [12] [13]. The metabolic implications of this lipid reprogramming are multifaceted, providing energy storage, membrane components for rapidly dividing cells, and precursors for signaling molecules that influence tumor progression. Similarly, kidney cancers develop a pronounced dependence on glutamine metabolism, utilizing this amino acid not only as a nitrogen donor for nucleotide synthesis but also as a carbon source for anaplerotic replenishment of TCA cycle intermediates [14] [15]. This metabolic flexibility enables tumors to adapt to nutrient-poor conditions and resist therapeutic interventions.

Beyond cancer cell-intrinsic metabolic changes, kidney tumors actively remodel their microenvironment through metabolic interactions that influence disease progression and treatment response [16]. The glycolytic TME becomes enriched in lactate and other metabolites that suppress immune cell function while promoting angiogenesis [17, 18]. Cancer-associated fibroblasts (CAFs) contribute to this metabolic symbiosis by providing alternative nutrient sources, while endothelial cells adapt to the hypoxic conditions by altering their own metabolic preferences [19, 20]. These complex interactions create therapeutic challenges but also reveal new vulnerabilities that could be exploited for more effective treatments.

The clinical implications of kidney cancer metabolism have become increasingly apparent with the development of targeted therapies. The recent FDA approval of belzutifan, a HIF-2 α inhibitor, validates the therapeutic potential of targeting cancer metabolism, while numerous other metabolic inhibitors are in clinical development [21]. However, significant challenges remain, including metabolic heterogeneity within tumors, the development of resistance mechanisms, and the need for reliable biomarkers to guide therapy selection. Emerging technologies such as metabolomic profiling, hyperpolarized MRI, and single-cell analysis are providing unprecedented insights into kidney cancer metabolism, enabling more precise targeting of metabolic vulnerabilities [22].

Metabolic pathways dysregulated in kidney cancer

Kidney cancer, particularly ccRCC undergoes significant metabolic reprogramming due to genetic and epigenetic alterations [11, 23]. A defining characteristic is the enhanced glycolytic flux, sustained even under normoxic conditions (the Warburg effect), driven by HIFs following loss of the VHL tumor suppressor (Figure 1) [24].

Lipid metabolism is also profoundly altered, with increased fatty acid uptake and storage to support membrane biosynthesis and energy reserves. Glutamine metabolism is similarly reconfigured, supplying critical precursors for nucleotide synthesis and glutathione production, thereby sustaining proliferation and redox balance [25]. Additionally, mitochondrial dysfunction impairs oxidative phosphorylation, further shifting dependence toward anaerobic metabolic pathways [26]. These adaptations not only fuel tumor growth and survival but also expose metabolic vulnerabilities that could be therapeutically targeted [27]. Elucidating these dysregulated pathways is essential for designing precision therapies to disrupt cancer metabolic dependencies and improve clinical outcomes.

Glycolysis and the warburg effect in kidney cancer

A hallmark of metabolic reprogramming in kidney cancer, particularly ccRCC, is the preferential utilization of glycolysis for energy production even in the presence of oxygen - a phenomenon termed the Warburg effect [6]. This metabolic shift is driven primarily by the constitutive stabilization of HIF-1 α and HIF-2 α due to loss of the VHL tumor suppressor. The enhanced glycolytic flux provides rapidly proliferating tumor cells with essential biosynthetic intermediates, including nucleotides, amino acids, and lipids, while simultaneously maintaining redox homeostasis through lactate production. Importantly, the Warburg effect supports tumor growth in the typically hypoxic microenvironment of renal carcinomas by reducing oxygen dependence for ATP generation [28]. This metabolic adaptation not only facilitates energy production but also creates a microenvironment that promotes immune evasion and therapeutic resistance [29]. The molecular underpinnings of glycolytic dysregulation in kidney cancer present promising targets for therapeutic intervention, including inhibitors of key glycolytic enzymes and HIF signaling pathways [30, 31].

Lipid metabolism reprogramming in kidney cancer

Kidney cancer exhibits profound alterations in lipid metabolism that support tumor growth and survival [32]. ccRCC is the most common renal malignancy, is particularly characterized by excessive lipid accumulation, visible histologically as cytoplasmic lipid droplets [33]. This metabolic rewiring is driven by multiple mechanisms, including HIF-mediated upregulation of lipid




Figure 1. Metabolic reprogramming drives kidney cancer progression. Stabilization and upregulation of HIF1 α and HIF2 α occur because of Loss of VHL. Metabolic pathways are dysregulated and rewired by alteration of several mediators (GLUT1, GLUT3, HK2, LDHA, etc.) which are activated by the action of HIF1 α and HIF2 α . Hypoxia is initiated in TME leading to immunosuppression through suppression of cytotoxic T-cells.

uptake receptors (e.g., CD36), enhanced de novo lipogenesis through increased expression of fatty acid synthase (FASN) and ATP-citrate lyase (ACLY), and impaired lipid oxidation due to mitochondrial dysfunction [13]. The resulting lipid-rich environment not only provides energy stores and membrane building blocks for rapidly proliferating tumor cells but also generates signaling molecules that promote tumor progression [34]. Notably, lipid droplets serve as reservoirs for cholesterol esters and phospholipids that can be mobilized to fuel cancer cell growth under nutrient-deprived conditions [35]. Furthermore, lipid-derived metabolites function as signaling molecules that modulate oncogenic pathways and contribute to the immunosuppressive TME. These metabolic adaptations present promising therapeutic targets, with several inhibitors of lipid metabolism currently under investigation for kidney cancer treatment.

Glutamine dependency and amino acid metabolism in kidney cancer

Renal cell carcinomas, particularly clear cell subtypes, demonstrate marked glutamine addiction as part of their metabolic reprogramming. This dependence stems from the tumor's need to replenish tricarboxylic acid (TCA) cycle intermediates (anaplerosis) and generate biosynthetic precursors for nucleotides, proteins, and antioxidants [36] [37]. The frequent loss of VHL and subsequent HIF stabilization upregulate glutamine transporters (ASCT2, SN2) and key enzymes like glutaminase (GLS), which converts glutamine to glutamate [38]. This metabolic adaptation becomes crucial in kidney cancer as mitochondrial dysfunction limits glucose-derived acetyl-CoA entry into the TCA cycle [6]. Beyond energy production, glutamine metabolism supports redox balance by maintaining glutathione levels and provides nitrogen for non-essential amino acid synthesis through transamination reactions. Interestingly, kidney tumors also alter other amino

acid pathways - notably upregulating serine/glycine metabolism for one-carbon units and modulating branched-chain amino acid catabolism. These interconnected amino acid fluxes create metabolic vulnerabilities, with preclinical studies showing sensitivity to glutaminase inhibitors and amino acid deprivation strategies. The emerging understanding of kidney cancer's amino acid metabolic network offers promising therapeutic avenues to target this nutrient dependency while potentially overcoming resistance to conventional therapies [31, 39].

Key drivers of metabolic reprogramming in kidney cancer

The metabolic rewiring observed in kidney cancer, specifically ccRCC is orchestrated by several interconnected molecular drivers [32]. The most prominent is the inactivation of the VHL tumor suppressor, which leads to constitutive stabilization of HIF-1 α and HIF-2 α [40]. These transcription factors activate a transcriptional program that upregulates glycolysis, enhances glutamine metabolism, and suppresses oxidative phosphorylation [41]. Concurrently, mutations in chromatin-modifying genes (e.g., PBRM1, SETD2) and activation of the PI3K-AKT-mTOR pathway further reshape metabolic networks by altering nutrient sensing and anabolic processes [42]. The TME, characterized by hypoxia and nutrient deprivation, exerts additional selective pressure that reinforces metabolic adaptations [43]. Importantly, these drivers converge to create a metabolic phenotype characterized by increased glucose and glutamine uptake, lipid droplet accumulation, and dependence on non-canonical nutrient utilization pathways. Understanding these key regulators provides critical insights for developing targeted therapies that disrupt cancer-specific metabolic dependencies while sparing normal tissues (**Figure 2**).

Hypoxia-inducible factors and their role in kidney cancer

pathways create a permissive metabolic environment characterized by heightened glucose and glutamine dependency, lipid droplet accumulation, and redox adaptation - all exploitable therapeutic vulnerabilities [64, 65]. Current targeted therapies (e.g., mTOR inhibitors, HIF-2 α antagonists) and emerging metabolic approaches aim to disrupt these oncogenic signaling-metabolism nexuses in kidney cancer.

Metabolic interactions in the tumor microenvironment of kidney cancer

The kidney cancer TME represents a complex metabolic ecosystem where neoplastic cells dynamically interact with stromal components, immune cells, and vasculature through nutrient competition and metabolic crosstalk [66]. Tumor cells preferentially utilize aerobic glycolysis (Warburg effect), creating lactate-rich niches that acidify the TME and suppress antitumor immune responses by impairing cytotoxic T-cell function while promoting regulatory T-cell (Treg) activity [67, 68]. CAFs contribute to this metabolic symbiosis by secreting amino acids, lipids, and lactate that fuel tumor growth through oxidative metabolism [69]. Endothelial cells adapt to the hypoxic conditions by upregulating angiogenic factors (VEGF, PDGF) in response to HIF stabilization, further perpetuating nutrient supply to rapidly proliferating tumor cells [70]. Importantly, tumor-infiltrating myeloid cells exhibit metabolic plasticity, shifting toward arginase-mediated immunosuppression in response to hypoxia and nutrient deprivation [71, 72]. These reciprocal metabolic interactions create a self-reinforcing protumorigenic milieu that facilitates immune evasion, therapeutic resistance, and metastatic progression. Emerging therapeutic strategies targeting these metabolic networks—such as lactate dehydrogenase inhibitors, glutamine antagonists, and immune-metabolic checkpoint modulators—aim to disrupt tumor-stromal co-dependencies and restore antitumor immunity in kidney cancer.

Crosstalk between tumor cells and stroma in kidney cancer

The bidirectional metabolic interplay between tumor cells and stromal components in kidney cancer creates a dynamic microenvironment that fuels disease progression. CAFs actively secrete lactate, pyruvate, and ketone bodies that tumor cells utilize as alternative energy substrates through oxidative phosphorylation, particularly under glucose-deprived conditions [69]. Conversely, tumor cells release glutamate and other oncometabolites that activate CAFs, inducing their transformation into myofibroblasts that further remodel the extracellular matrix [73]. This metabolic symbiosis extends to endothelial cells, where HIF-driven VEGF secretion from tumor cells promotes angiogenesis, while the resulting neovasculature provides nutrients and oxygen that sustain tumor growth [74]. Adipocytes in perirenal fat deposits contribute free fatty acids that tumor cells internalize through CD36-mediated uptake, supporting membrane biosynthesis and energy storage [75]. Importantly, this crosstalk is mediated by exosomal transfer of miRNAs and metabolic enzymes that reprogram recipient cells. The resulting metabolic coupling not only enhances tumor survival under stress conditions but also creates therapeutic resistance by establishing redundant nutrient acquisition pathways. Targeting these tumor-stroma metabolic interactions – through approaches like CAF depletion, anti-angiogenic therapy, or lipid metabolism inhibition – represents a promising strategy to disrupt the tumor-supportive niche in kidney cancer.

Immune cell metabolism and immunosuppression in the kidney cancer microenvironment

The metabolic landscape of kidney cancer actively shapes antitumor immunity by imposing nutrient constraints and altering immune cell functionality within the TME. Tumor cells outcompete infiltrating lymphocytes for glucose through elevated expression of GLUT1 and hexokinase-2, forcing cytotoxic T cells into a hypofunctional state characterized by impaired glycolysis and reduced interferon- γ production [76]. Conversely, myeloid-derived suppressor cells (MDSCs) and regulatory T cells (Tregs) thrive in this metabolically hostile environment by preferentially utilizing fatty acid oxidation and oxidative phosphorylation, which support their immunosuppressive functions [77]. The accumulation of tumor-derived lactate and kynurenine further reinforces immunosuppression by inhibiting natural killer cell activity while promoting the polarization of tumor-associated macrophages toward an M2 phenotype. Notably, the hypoxic tumor core drives PD-L1 upregulation on both cancer cells and infiltrating myeloid cells through HIF-1 α stabilization, creating an immune checkpoint-rich environment (**Figure 3**) [78]. These metabolic constraints contribute to the limited efficacy of immunotherapies in kidney cancer, prompting investigations into metabolic modulators – such as lactate dehydrogenase inhibitors and IDO1 antagonists – that may reverse immunosuppression and enhance checkpoint blockade responses [79]. Understanding these immunometabolic interactions provides critical insights for developing combination strategies that simultaneously target tumor metabolism and immune evasion mechanisms.

Angiogenesis and nutrient supply in kidney cancer progression

Kidney cancer orchestrates a robust angiogenic response to sustain its metabolic demands through complex interactions between tumor cells and the vascular microenvironment. The characteristic VHL/HIF axis activation in ccRCC drives excessive vascular endothelial growth factor (VEGF) production [53], stimulating the formation of disorganized, hyperpermeable tumor vasculature. These aberrant vessels, while providing increased nutrient and oxygen supply, create a paradoxical state of chronic hypoxia due to their structural abnormalities and inefficient perfusion. Tumor cells adapt by further upregulating HIF-dependent glycolytic enzymes and glucose transporters, establishing a self-perpetuating cycle of metabolic demand and vascular recruitment [80, 81]. The resulting vasculature not only delivers glucose and glutamine but also serves as a conduit for lipid uptake from circulating lipoproteins, supporting the lipid droplet accumulation characteristic of ccRCC. Importantly, the angiogenic switch enables metastatic dissemination by providing tumor cells access to systemic circulation while simultaneously creating an immunosuppressive microenvironment through VEGF-mediated inhibition of dendritic cell maturation [82]. This understanding has led to the clinical success of anti-angiogenic therapies, though their efficacy is often limited by the emergence of alternative nutrient acquisition strategies, including enhanced macropinocytosis and vascular co-option. Current research focuses on combining VEGF pathway inhibitors with metabolic or immunotherapeutic agents to more effectively starve tumors while preventing compensatory adaptations.

Diagnostic and prognostic implications of metabolic alterations in kidney cancer

The distinct metabolic profile of kidney cancer offers clinically valuable biomarkers for disease detection, stratification, and monitoring. The hallmark lipid and glycogen accumulation in ccRCC provides diagnostic utility, with imaging modalities like chemical-shift MRI effectively distinguishing malignant lesions from benign renal masses by detecting intracellular lipid content




Figure 3. Immunosuppressive molecular mechanism in kidney cancer. LDH facilitates conversion of tumor-derived Lactate from pyruvate. The accumulation of LDH acidifies the TME leading to inhibition of cytotoxic T-cell functions, NK cell functions, and modulation of tumor associated macrophages (TAM). As a result, Treg cells are activated which further promotes cancer cell progression.

[11]. Metabolic alterations also carry prognostic significance—elevated FDG-PET avidity correlates with aggressive tumor behavior [83], while high expression of glycolytic enzymes (HK2, LDHA) and glutamine transporters (ASCT2) predicts poor survival outcomes [84]. Liquid biopsy approaches are increasingly detecting circulating tumor metabolites (succinate, 2-hydroxyglutarate) that reflect underlying mutations in IDH genes, enabling non-invasive molecular classification [85, 86]. Notably, the extent of metabolic rewiring mirrors disease progression, with metastatic lesions demonstrating amplified HIF activation and more pronounced Warburg effect compared to primary tumors. Emerging metabolic signatures, such as the ratio of ketone bodies to free fatty acids in serum, show promise for monitoring therapeutic response and detecting early recurrence. These metabolic readouts not only improve clinical decision-making but also reveal actionable targets, as tumors with specific metabolic vulnerabilities (e.g., glutamine dependency or defective oxidative phosphorylation) may show preferential sensitivity to pathway inhibitors. The integration of metabolic profiling with conventional imaging and genomic data is paving the way for precision oncology approaches in kidney cancer management.

Metabolic biomarkers in kidney cancer: current applications and emerging potential

The unique metabolic rewiring of kidney cancer has yielded clinically relevant biomarkers that enhance diagnostic precision, prognostic stratification, and therapeutic monitoring. ccRCC-specific metabolic signatures—including elevated circulating succinate levels from pseudohypoxic drive and increased urinary N-acetylaspartate reflecting altered lipid metabolism—provide non-invasive diagnostic indicators that complement imaging findings [87]. Prognostically, immunohistochemical detection of key metabolic enzymes (CAIX, GLUT1) in tumor tissues stratifies patient risk, while liquid biopsy profiles measuring kynurenine/tryptophan ratios or branched-chain amino acid

patterns predict immunotherapy response [88]. Advanced imaging biomarkers, particularly 18F-FDG PET avidity and hyperpolarized 13C-pyruvate MRI, quantitatively map tumor glycolytic activity, correlating with tumor grade and metastatic potential [89]. Emerging mass spectrometry-based metabolomics now identify signature perturbations in TCA cycle intermediates (fumarate, 2-HG) that reveal underlying genetic alterations (FH/SDH mutations) and guide targeted therapy selection [90]. Notably, dynamic changes in serum acylcarnitine profiles and extracellular vesicle-derived metabolic enzymes show promise for real-time treatment monitoring. These biomarkers collectively address critical clinical challenges in kidney cancer management, from differentiating indolent from aggressive disease to detecting micro-metastases and overcoming therapeutic resistance. Their integration into multi-omics diagnostic platforms is advancing personalized management strategies that align tumor-specific metabolic vulnerabilities with precision therapies.

Imaging techniques for metabolic profiling in kidney cancer

Advanced imaging modalities now enable non-invasive metabolic profiling of kidney tumors, providing critical diagnostic and prognostic information while guiding treatment decisions. Positron emission tomography (PET) using 18F-fluorodeoxyglucose (FDG) remains the cornerstone for evaluating glycolytic activity, with standardized uptake values (SUVmax) correlating with tumor aggressiveness and metastatic potential [91, 92]. Novel PET tracers targeting other metabolic pathways—such as 11C-acetate for lipid metabolism and 18F-fluoroglutamine for amino acid uptake—are expanding the metabolic profiling capabilities [93]. Magnetic resonance spectroscopy (MRS) offers complementary data by quantifying endogenous metabolites, including elevated choline peaks reflecting membrane turnover and reduced citrate levels characteristic of ccRCC. Emerging hyperpolarized 13C-pyruvate MRI techniques dynamically track real-time conversion of pyruvate to lactate, directly visualizing Warburg effect activity

Table 1. Anticancer drugs effective in kidney cancer.

Name of the drug	Manufacturer	Mechanism	Reference
Lonidamine	Angelini pharma	Hexokinase-2 inhibitor	[102]
Belzutifan	Merck & Co.	Targeting HIF-2 α	[103]
Telaglenastat	Calithera biosciences	Glutamine pathway inhibitor	[104]
TVB-2640	Sagimet biosciences	FASN inhibitor	[105]
FX11	Albert Einstein college of medicine	LDHA inhibitor	[106]

with unprecedented spatial resolution [94]. Chemical shift imaging reliably detects intracellular lipid content, distinguishing clear cell from non-clear cell variants with >90% accuracy [95]. These functional imaging approaches are being integrated with radiomic analysis of conventional CT/MRI to create multiparametric metabolic signatures that predict treatment response and monitor therapeutic efficacy. The non-invasive nature of metabolic imaging positions it as an ideal tool for serial assessment during therapy, particularly for evaluating emerging metabolism-targeted treatments like HIF-2 α inhibitors and glutaminase blockers.

Therapeutic targeting of metabolic pathways in kidney cancer

The distinct metabolic dependencies of kidney cancer present promising opportunities for targeted therapeutic intervention (Table 1). Current strategies focus on disrupting the glycolytic flux through inhibitors of rate-limiting enzymes such as hexokinase-2 (lonidamine) or lactate dehydrogenase (FX11), which preferentially affect tumor cells exhibiting the Warburg effect [96]. The pivotal role of HIF-2 α in ccRCC metabolism has been successfully targeted by belzutifan, an FDA-approved inhibitor that attenuates pseudohypoxic signaling and its downstream metabolic effects [97]. Simultaneously, glutamine pathway inhibitors like CB-839 (telaglenastat) exploit the tumor's reliance on anaplerosis by blocking glutaminase-mediated conversion to glutamate [27]. Emerging approaches target lipid metabolism through FASN inhibitors (TVB-2640) or disrupt redox balance by inhibiting NAD⁺ biosynthesis [98]. Notably, these metabolic therapies demonstrate synergistic potential when combined with existing anti-angiogenics or immunotherapies, as evidenced by enhanced T-cell infiltration following lactate export blockade [99]. Second-generation strategies now explore TME-specific targets, including acidosis-neutralizing agents and macrophage-directed metabolic modulators [100]. The development of pharmacodynamic biomarkers—such as hyperpolarized MRI-detected pyruvate-to-lactate conversion rates—enables real-time monitoring of metabolic drug effects, facilitating personalized treatment optimization [101]. This multifaceted approach to metabolic targeting addresses both cancer cell-intrinsic dependencies and tumor-extrinsic metabolic crosstalk, offering new avenues to overcome therapeutic resistance in kidney cancer.

Inhibitors of glycolysis and HIF signaling in kidney cancer therapeutics

The targeting of glycolytic and HIF signaling pathways represents a precision medicine approach for kidney cancer, capitalizing on the tumor's hallmark metabolic vulnerabilities. HIF-2 α antagonists such as belzutifan (MK-6482) have demonstrated

clinical efficacy by specifically disrupting the pseudohypoxic transcriptional program in VHL-deficient tumors, reducing expression of glycolytic enzymes (HK2, LDHA) and glucose transporters (GLUT1/3) [107]. Parallel strategies employ small molecule inhibitors of rate-limiting glycolytic components—including 2-deoxyglucose (glycolytic inhibitor) and PFK158 (PFKFB3 blocker)—to starve tumors of their preferred energy source while sparing normal cells that retain oxidative phosphorylation capacity [108]. Particularly promising are dual-action compounds that concurrently target HIF signaling and glycolysis, such as PT2385 derivatives that destabilize HIF-2 α while inhibiting hexokinase activity [109]. These approaches show synergistic potential when combined with anti-angiogenic therapies, as HIF inhibition normalizes tumor vasculature while glycolytic blockade prevents metabolic adaptation. Resistance mechanisms, including upregulation of alternate HIF isoforms or activation of compensatory nutrient salvage pathways, are being addressed through next-generation inhibitors with improved target specificity and combination regimens incorporating glutaminase blockers. The development of PET-based biomarkers (18F-FDG, 18F-fluoromisonidazole) enables real-time monitoring of therapeutic response, facilitating dose optimization for these metabolism-targeted agents [110]. This therapeutic paradigm exemplifies how understanding cancer-specific metabolic dependencies can yield targeted treatments with potentially fewer off-target effects than conventional therapies.

Targeting lipid and amino acid metabolism in kidney cancer therapy

Emerging therapeutic strategies are exploiting the deregulated lipid and amino acid metabolism that underlies kidney cancer pathogenesis. The characteristic lipid droplet accumulation in clear cell RCC has prompted development of fatty acid synthase (FASN) inhibitors like TVB-2640, which disrupt de novo lipogenesis and induce tumor-specific apoptosis by depriving cancer cells of membrane precursors and signaling lipids [111]. Concurrently, inhibitors of sterol regulatory element-binding proteins (SREBP) such as fatostatin are being evaluated to block the lipogenic transcription program driven by HIF and mTOR pathways [112]. In amino acid metabolism, glutaminase inhibitors (telaglenastat) and ASCT2 blockers (V-9302) are showing promise in clinical trials by restricting tumor access to glutamine—a crucial nitrogen and carbon source for ccRCC proliferation [113]. Notably, these approaches synergize with existing therapies: lipid metabolism inhibitors enhance anti-angiogenic efficacy by reducing VEGF production, while amino acid restriction potentiates immunotherapy by alleviating immunosuppressive tryptophan/kynurenine pathways. Advanced patient stratification

using lipidomic profiles and PET imaging with glutamine analogs (18F-FGln) is enabling precision targeting of these metabolic vulnerabilities. The simultaneous targeting of both lipid and amino acid pathways represents a multipronged strategy to overwhelm tumor metabolic plasticity and overcome treatment resistance in kidney cancer.

Combination therapies: metabolic drugs with immunotherapy/TKI in kidney cancer treatment

The strategic integration of metabolic modulators with immunotherapy and tyrosine kinase inhibitors (TKIs) represents a paradigm shift in kidney cancer treatment, addressing both tumor-intrinsic vulnerabilities and microenvironmental immunosuppression. Preclinical studies demonstrate that HIF-2 α inhibitors (belzutifan) synergize with PD-1/PD-L1 blockade by alleviating hypoxia-driven immunosuppression while normalizing aberrant tumor vasculature when combined with VEGF-targeted TKIs [114]. Clinically, glutaminase inhibitors (telaglenastat) are being evaluated with pembrolizumab to simultaneously restrict tumor bioenergetics and enhance T-cell function by reducing myeloid-derived suppressor cell (MDSC) accumulation in the TME [115]. Similarly, lactate dehydrogenase inhibitors (GSK2837808A) are showing promise in combination regimens by reversing the lactate-mediated suppression of cytotoxic T lymphocytes while maintaining anti-angiogenic effects of TKIs [116]. Emerging trial data reveal that these combinations yield durable responses by targeting complementary resistance mechanisms—metabolic drugs prevent the glycolytic adaptation that often limits TKI efficacy, while immunotherapy counters the immunosuppressive effects of metabolic stress. Advanced biomarker strategies, including metabolic PET imaging (18F-FDG, 18F-FSPG) and immune-metabolic profiling of tumor biopsies, are enabling real-time monitoring of these synergistic effects [117]. This multidimensional therapeutic approach capitalizes on the interconnected nature of metabolic and signaling networks in kidney cancer, offering new avenues to overcome treatment resistance and improve long-term outcomes.

Challenges and future perspectives in targeting kidney cancer metabolism

Despite significant advances in understanding metabolic reprogramming in kidney cancer, several challenges hinder the clinical translation of metabolism-targeted therapies. Tumor heterogeneity and metabolic plasticity enable cancer cells to rapidly switch between energy pathways, fostering resistance to single-agent therapies that target specific metabolic nodes [118]. The dual role of certain metabolites—such as lactate functioning as both a fuel source and immunosuppressive agent—complicates therapeutic interventions [119], while systemic toxicity remains a concern when inhibiting fundamental metabolic processes shared by normal cells. Current limitations in real-time metabolic imaging and biomarker validation further impede personalized treatment strategies. Future directions include the development of multi-target inhibitors that simultaneously block compensatory pathways, along with advanced drug delivery systems like nanoparticle conjugates to enhance tumor specificity. Artificial intelligence-driven analysis of multi-omics data promises to uncover novel metabolic vulnerabilities and optimize combination regimens [120]. Clinically, the integration of metabolic modulators with immunotherapy and targeted agents in rationally designed trials—guided by robust pharmacodynamic biomarkers—will be critical. Additionally, exploring circadian regulation of cancer metabolism and host-microbiome metabolic interactions may reveal unexpected therapeutic opportunities. Overcoming these

challenges requires a multidisciplinary approach that bridges basic metabolic research with innovative clinical trial designs, ultimately paving the way for more effective, durable treatments in kidney cancer.

Resistance to metabolic therapies in kidney cancer

The emergence of resistance to metabolism-targeted agents in kidney cancer stems from the remarkable metabolic plasticity and genetic adaptability of tumor cells. A primary mechanism involves compensatory upregulation of alternative nutrient acquisition pathways—for instance, tumors treated with glutaminase inhibitors frequently activate macropinocytosis to scavenge extracellular proteins or amplify ASCT2-independent glutamine transport systems [121]. Similarly, glycolytic blockade often triggers a metabolic shift toward oxidative phosphorylation through mitochondrial genome amplification or increased fatty acid β -oxidation [122]. Epigenetic remodeling enables rapid adaptation, with demethylation of metabolic gene promoters facilitating expression of bypass pathways under therapeutic pressure. The TME further contributes to resistance through metabolic symbiosis, where stromal cells supply metabolites (lactate, ketones) that rescue treated tumor cells from energy crisis. Heterogeneous expression of metabolic enzymes across tumor subclones creates inherent resistance reservoirs, while HIF stabilization in perinecrotic regions maintains tumor survival despite therapy [123]. Emerging strategies to overcome resistance include intermittent dosing to prevent adaptive responses, dual targeting of complementary metabolic nodes (e.g., concurrent glycolysis and OXPHOS inhibition), and combining metabolic drugs with epigenetic modifiers to limit transcriptional adaptation. The development of functional metabolic imaging techniques (hyperpolarized MRI, metabolic PET tracers) now enables real-time monitoring of these resistance mechanisms, guiding adaptive therapeutic strategies in clinical trials.

Emerging technologies and novel therapeutic targets in kidney cancer metabolism

Recent advances in multi-omics technologies and high-resolution metabolic imaging are uncovering novel therapeutic vulnerabilities in kidney cancer metabolism. Single-cell metabolomics has revealed previously unappreciated metabolic heterogeneity within tumors, identifying rare subpopulations with dependencies on cysteine or one-carbon metabolism that could be targeted with new small-molecule inhibitors [124]. CRISPR-based metabolic gene screening has pinpointed hexosamine biosynthesis and serine/glycine conversion as essential pathways in VHL-deficient cells, while spatial transcriptomics maps metabolic crosstalk between tumor and immune cells within the TME [125]. Emerging therapeutic targets include the mitochondrial pyruvate carrier (MPC) – inhibition of which selectively starves kidney cancer cells of TCA cycle intermediates – and the cystine/glutamate antiporter xCT, which maintains redox balance in metastatic lesions. Nanotechnology approaches are enabling targeted delivery of metabolic drugs, such as nanoparticle-encapsulated glutaminase inhibitors that preferentially accumulate in tumors. Meanwhile, AI-driven analysis of metabolic flux data is predicting patient-specific vulnerabilities by modeling individual tumor metabolic networks. These innovations are being translated clinically through innovative trial designs, including basket trials testing metabolic therapies based on molecular features rather than histology, and window-of-opportunity studies using hyperpolarized ^{13}C -MRI to quantify real-time drug effects on tumor metabolism [101]. Together, these technological advances are expanding the arsenal of metabolism-targeted therapies while enabling precision

approaches tailored to individual patient tumors.

Conclusion

The metabolic reprogramming of kidney cancer represents a fundamental hallmark of the disease, driven by genetic alterations, microenvironmental pressures, and adaptive survival mechanisms. Key findings highlight the central role of HIF-mediated pseudohypoxia, dysregulated lipid and amino acid metabolism, and bidirectional crosstalk between tumor and stromal cells in promoting tumor progression and therapeutic resistance. While significant progress has been made in targeting these pathways—exemplified by the clinical success of HIF-2 α inhibitors—challenges such as metabolic plasticity and immunosuppressive niche formation persist. Future directions will require innovative approaches, including the integration of multi-omics technologies for patient stratification, development of dual-targeting metabolic agents, and rational combinations with immunotherapy to address tumor heterogeneity. Advances in metabolic imaging and AI-driven biomarker discovery are poised to accelerate precision medicine strategies, enabling real-time monitoring of treatment efficacy. As our understanding of kidney cancer metabolism evolves, so too will opportunities to develop more effective therapies that disrupt metabolic vulnerabilities while minimizing systemic toxicity. Ultimately, translating these insights into clinical practice demands collaborative efforts between basic researchers, clinicians, and bioengineers to overcome current limitations and improve outcomes for patients with kidney cancer.

Acknowledgements

None.

Ethical policy

Non applicable.

Availability of data and materials

All data generated or analysed during this study are included in this publication.

Author contributions

Elena Tena Edo contributed to design of the work, data collection, and drafting the article.

Competing interests

The author declares no competing interests.

Funding

None.

References

- Hu J, Wang S-G, Hou Y, Chen Z, Liu L, Li R, Li N, Zhou L, Yang Y, Wang L et al: Multi-omic profiling of clear cell renal cell carcinoma identifies metabolic reprogramming associated with disease progression. *Nature Genetics* 2024, 56(3): 442-457.
- Sudarshan S, Karam JA, Brugarolas J, Thompson RH, Uzzo R, Rini B, Margulis V, Patard JJ, Escudier B, Linehan WM: Metabolism of kidney cancer: from the lab to clinical practice. *Eur Urol* 2013, 63(2): 244-251.
- Carthew RW: Gene Regulation and Cellular Metabolism: An Essential Partnership. *Trends Genet* 2021, 37(4): 389-400.
- Liu Y, Zhao Y, Song H, Li Y, Liu Z, Ye Z, Zhao J, Wu Y, Tang J, Yao M: Metabolic reprogramming in tumor immune microenvironment: Impact on immune cell function and therapeutic implications. *Cancer Letters* 2024, 597: 217076.
- Mao Y, Xia Z, Xia W, Jiang P: Metabolic reprogramming, sensing, and cancer therapy. *Cell Reports* 2024, 43(12): 115064.
- Chen Z, Zhang X: The role of metabolic reprogramming in kidney cancer. *Front Oncol* 2024, 14: 1402351.
- Haase VH: The VHL/HIF oxygen-sensing pathway and its relevance to kidney disease. *Kidney International* 2006, 69(8): 1302-1307.
- Zhang J, Yao M, Xia S, Zeng F, Liu Q: Systematic and comprehensive insights into HIF-1 stabilization under normoxic conditions: implications for cellular adaptation and therapeutic strategies in cancer. *Cellular & Molecular Biology Letters* 2025, 30(1): 2.
- Nsiah NY, Morgan AB, Donkor N, Inman DM: Long-term HIF-1 α stabilization reduces respiration, promotes mitophagy, and results in retinal cell death. *Scientific Reports* 2023, 13(1): 20541.
- Mathew M, Nguyen NT, Bhutia YD, Sivaprakasam S, Ganapathy V: Metabolic Signature of Warburg Effect in Cancer: An Effective and Obligatory Interplay between Nutrient Transporters and Catabolic/Anabolic Pathways to Promote Tumor Growth. *Cancers (Basel)* 2024, 16(3): 504.
- Sanchez DJ, Simon MC: Genetic and metabolic hallmarks of clear cell renal cell carcinoma. *Biochim Biophys Acta Rev Cancer* 2018, 1870(1): 23-31.
- Wang J, Guo H, Zheng L-F, Li P, Zhao T-J: Context-specific fatty acid uptake is a finely-tuned multi-level effort. *Trends in Endocrinology & Metabolism* 2024, <https://doi.org/https://doi.org/10.1016/j.tem.2024.10.001>. Epub ahead of print.
- Schroeder B, Vander Steen T, Espinoza I, Venkatapoorana CMK, Hu Z, Silva FM, Regan K, Cuyàs E, Meng XW, Verdura S et al: Fatty acid synthase (FASN) regulates the mitochondrial priming of cancer cells. *Cell Death & Disease* 2021, 12(11): 977.
- Abu Aboud O, Habib SL, Trott J, Stewart B, Liang S, Chaudhari AJ, Sutcliffe J, Weiss RH: Glutamine Addition in Kidney Cancer Suppresses Oxidative Stress and Can Be Exploited for Real-Time Imaging. *Cancer Res* 2017, 77(23): 6746-6758.
- Jin J, Byun J-K, Choi Y-K, Park K-G: Targeting glutamine metabolism as a therapeutic strategy for cancer. *Experimental & Molecular Medicine* 2023, 55(4): 706-715.
- Elia I, Haigis MC: Metabolites and the tumour microenvironment: from cellular mechanisms to systemic metabolism. *Nat Metab* 2021, 3(1): 21-32.
- Wang Z-H, Peng W-B, Zhang P, Yang X-P, Zhou Q: Lactate in the tumour microenvironment: From immune modulation to therapy. *EBioMedicine* 2021, 73: 103627.
- Pérez-Tomás R, Pérez-Guillén I: Lactate in the Tumor Microenvironment: An Essential Molecule in Cancer Progression and Treatment. *Cancers (Basel)* 2020, 12(11): 3244.
- Li Z, Sun C, Qin Z: Metabolic reprogramming of cancer-associated fibroblasts and its effect on cancer cell reprogramming. *Theranostics* 2021, 11(17): 8322-8336.
- Xia B, Qiu L, Yue J, Si J, Zhang H: The metabolic crosstalk of cancer-associated fibroblasts and tumor cells: Recent advances and future perspectives. *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer* 2024, 1879(6): 189190.
- Wu X, Lazris D, Wong R, Tykodi SS: Belzutifan for the treatment of renal cell carcinoma. *Ther Adv Med Oncol* 2025, 17: 17588359251317846.
- Jonasch E, Donskov F, Iliopoulos O, Rathmell WK, Narayan VK, Maughan BL, Oudard S, Else T, Maranchie JK, Welsh SJ et al: Belzutifan for Renal Cell Carcinoma in von Hippel-Lindau Disease. *N Engl J Med* 2021, 385(22): 2036-2046.

23. Zheng X, Liu Y, Yang Z, Tian Y: Metabolic reprogramming and immune microenvironment profiling in clear cell renal cell carcinoma: implications for prognosis, targeted therapy, and drug resistance. *Discover Oncology* 2025, 16(1): 850.
24. Kierans SJ, Taylor CT: Regulation of glycolysis by the hypoxia-inducible factor (HIF): implications for cellular physiology. *The Journal of Physiology* 2021, 599(1): 23-37.
25. Yoo HC, Yu YC, Sung Y, Han JM: Glutamine reliance in cell metabolism. *Experimental & Molecular Medicine* 2020, 52(9): 1496-1516.
26. Bhatti JS, Bhatti GK, Reddy PH: Mitochondrial dysfunction and oxidative stress in metabolic disorders — A step towards mitochondria based therapeutic strategies. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* 2017, 1863(5): 1066-1077.
27. Tufail M, Jiang C-H, Li N: Altered metabolism in cancer: insights into energy pathways and therapeutic targets. *Molecular Cancer* 2024, 23(1): 203.
28. Liberti MV, Locasale JW: The Warburg Effect: How Does it Benefit Cancer Cells? *Trends Biochem Sci* 2016, 41(3): 211-218.
29. Zhang H, Li S, Wang D, Liu S, Xiao T, Gu W, Yang H, Wang H, Yang M, Chen P: Metabolic reprogramming and immune evasion: the interplay in the tumor microenvironment. *Biomarker Research* 2024, 12(1): 96.
30. Rabelink TJ, Carmeliet P: Glycolytic adaptation and progression of kidney disease. *Nature Reviews Nephrology* 2018, 14(2): 75-76.
31. Rathmell WK, Rathmell JC, Linehan WM: Metabolic Pathways in Kidney Cancer: Current Therapies and Future Directions. *J Clin Oncol* 2018, <https://doi.org/10.1200/jco.2018.79.2309>. Epub ahead of print.:Jco2018792309.
32. Zhang Y, Zhang S, Sun H, Xu L: The pathogenesis and therapeutic implications of metabolic reprogramming in renal cell carcinoma. *Cell Death Discovery* 2025, 11(1): 186.
33. Qi X, Li Q, Che X, Wang Q, Wu G: The Uniqueness of Clear Cell Renal Cell Carcinoma: Summary of the Process and Abnormality of Glucose Metabolism and Lipid Metabolism in ccRCC. *Front Oncol* 2021, 11: 727778.
34. Corn KC, Windham MA, Rafat M: Lipids in the tumor microenvironment: From cancer progression to treatment. *Progress in Lipid Research* 2020, 80: 101055.
35. Olzmann JA, Carvalho P: Dynamics and functions of lipid droplets. *Nature Reviews Molecular Cell Biology* 2019, 20(3): 137-155.
36. Eniafe J, Jiang S: The functional roles of TCA cycle metabolites in cancer. *Oncogene* 2021, 40(19): 3351-3363.
37. Anderson NM, Mucka P, Kern JG, Feng H: The emerging role and targetability of the TCA cycle in cancer metabolism. *Protein Cell* 2018, 9(2): 216-237.
38. Haase VH: The VHL tumor suppressor: master regulator of HIF. *Curr Pharm Des* 2009, 15(33): 3895-3903.
39. Do LK, Lee HM, Ha Y-S, Lee C-H, Kim J: Amino acids in cancer: Understanding metabolic plasticity and divergence for better therapeutic approaches. *Cell Reports* 2025, 44(4): 115529.
40. Miller F, Kentsis A, Osman R, Pan Z-Q: Inactivation of VHL by Tumorigenic Mutations That Disrupt Dynamic Coupling of the pVHL-Hypoxia-inducible Transcription Factor-1 α Complex. *Journal of Biological Chemistry* 2005, 280(9): 7985-7996.
41. Ni X, Lu C-p, Xu G-q, Ma J-j: Transcriptional regulation and post-translational modifications in the glycolytic pathway for targeted cancer therapy. *Acta Pharmacologica Sinica* 2024, 45(8): 1533-1555.
42. Walton J, Lawson K, Prinos P, Finelli A, Arrowsmith C, Ailles L: PBRM1, SETD2 and BAP1 — the trinity of 3p in clear cell renal cell carcinoma. *Nature Reviews Urology* 2023, 20(2): 96-115.
43. Singleton DC, Macann A, Wilson WR: Therapeutic targeting of the hypoxic tumour microenvironment. *Nature Reviews Clinical Oncology* 2021, 18(12): 751-772.
44. Hoefflin R, Harlander S, Schäfer S, Metzger P, Kuo F, Schönenberger D, Adlesic M, Peighambari A, Seidel P, Chen C-y et al: HIF-1 α and HIF-2 α differently regulate tumour development and inflammation of clear cell renal cell carcinoma in mice. *Nature Communications* 2020, 11(1): 4111.
45. Tanimoto K, Makino Y, Pereira T, Poellinger L: Mechanism of regulation of the hypoxia-inducible factor-1 alpha by the von Hippel-Lindau tumor suppressor protein. *Embo j* 2000, 19(16): 4298-4309.
46. Hayashi Y, Yokota A, Harada H, Huang G: Hypoxia/pseudohypoxia-mediated activation of hypoxia-inducible factor-1 α in cancer. *Cancer Sci* 2019, 110(5): 1510-1517.
47. Yang Z, Wei X, Ji C, Ren X, Su W, Wang Y, Zhou J, Zhao Z, Zhou P, Zhao K et al: OGT/HIF--2 α axis promotes the progression of clear cell renal cell carcinoma and regulates its sensitivity to ferroptosis. *iScience* 2023, 26(11): 108148.
48. Sun RC, Denko NC: Hypoxic regulation of glutamine metabolism through HIF1 and SIAH2 supports lipid synthesis that is necessary for tumor growth. *Cell Metab* 2014, 19(2): 285-292.
49. Cowman SJ, Koh MY: Revisiting the HIF switch in the tumor and its immune microenvironment. *Trends in Cancer* 2022, 8(1): 28-42.
50. Shenoy N, Pagliaro L: Sequential pathogenesis of metastatic VHL mutant clear cell renal cell carcinoma: putting it together with a translational perspective. *Annals of Oncology* 2016, 27(9): 1685-1695.
51. Roldan-Romero JM, Beuselinck B, Santos M, Rodriguez-Moreno JF, Lanillos J, Calsina B, Gutierrez A, Tang K, Lainez N, Puente J et al: PTEN expression and mutations in TSC1, TSC2 and MTOR are associated with response to rapalogs in patients with renal cell carcinoma. *Int J Cancer* 2020, 146(5): 1435-1444.
52. Hassan B, Akcakanat A, Holder AM, Meric-Bernstam F: Targeting the PI3-kinase/Akt/mTOR signaling pathway. *Surg Oncol Clin N Am* 2013, 22(4): 641-664.
53. Mazumder S, Higgins PJ, Samarakoon R: Downstream Targets of VHL/HIF- α Signaling in Renal Clear Cell Carcinoma Progression: Mechanisms and Therapeutic Relevance. *Cancers (Basel)* 2023, 15(4): 1316.
54. Yoo A, Tang C, Zucker M, Fitzgerald K, DiNatale RG, Rappold PM, Weiss K, Freeman B, Lee C-H, Schultz N et al: Genomic and Metabolic Hallmarks of SDH- and FH-deficient Renal Cell Carcinomas. *European Urology Focus* 2022, 8(5): 1278-1288.
55. Yuan X, Ruan W, Bobrow B, Carmeliet P, Eltzschig HK: Targeting hypoxia-inducible factors: therapeutic opportunities and challenges. *Nature Reviews Drug Discovery* 2024, 23(3): 175-200.
56. Xie D, Li G, Zheng Z, Zhang X, Wang S, Jiang B, Li X, Wang X, Wu G: The molecular code of kidney cancer: A path of discovery for gene mutation and precision therapy. *Molecular Aspects of Medicine* 2025, 101: 101335.
57. Aoki M, Fujishita T: Oncogenic Roles of the PI3K/AKT/mTOR Axis. *Curr Top Microbiol Immunol* 2017, 407: 153-189.
58. Glaviano A, Foo ASC, Lam HY, Yap KCH, Jacot W, Jones RH, Eng H, Nair MG, Makvandi P, Goerger B et al: PI3K/AKT/mTOR signaling transduction pathway and targeted therapies in cancer. *Mol Cancer* 2023, 22(1): 138.
59. Kapitsinou PP, Haase VH: The VHL tumor suppressor and HIF: insights from genetic studies in mice. *Cell Death & Differentiation* 2008, 15(4): 650-659.
60. Stefan E, Bister K: MYC and RAF: Key Effectors in Cellular Signaling and Major Drivers in Human Cancer. *Curr Top Microbiol Immunol* 2017, 407: 117-151.
61. Ritt DA, Abreu-Blanco MT, Bindu L, Durrant DE, Zhou M, Specht SI, Stephen AG, Holderfield M, Morrison DK: Inhibition of Ras/Raf/MEK/ERK Pathway Signaling by a Stress-Induced Phospho-Regulatory Circuit. *Molecular Cell* 2016, 64(5): 875-887.
62. Sakamoto T, Weng JS, Hara T, Yoshino S, Kozuka-Hata H, Oyama M, Seiki M: Hypoxia-inducible factor 1 regulation through cross

- talk between mTOR and MT1-MMP. *Mol Cell Biol* 2014, 34(1): 30-42.
63. Liu J, Zhang C, Hu W, Feng Z: Tumor suppressor p53 and metabolism. *J Mol Cell Biol* 2019, 11(4): 284-292.
 64. Tufail M, Jiang CH, Li N: Altered metabolism in cancer: insights into energy pathways and therapeutic targets. *Mol Cancer* 2024, 23(1): 203.
 65. Bosch M, Sweet MJ, Parton RG, Pol A: Lipid droplets and the host-pathogen dynamic: FAtal attraction? *J Cell Biol* 2021, 220(8): e202104005.
 66. Kay EJ, Zanivan S: The tumor microenvironment is an ecosystem sustained by metabolic interactions. *Cell Reports* 2025, 44(3): 115432.
 67. Zhou D, Duan Z, Li Z, Ge F, Wei R, Kong L: The significance of glycolysis in tumor progression and its relationship with the tumor microenvironment. *Front Pharmacol* 2022, 13: 1091779.
 68. Zhang A, Fan T, Liu Y, Yu G, Li C, Jiang Z: Regulatory T cells in immune checkpoint blockade antitumor therapy. *Molecular Cancer* 2024, 23(1): 251.
 69. Avagliano A, Granato G, Ruocco MR, Romano V, Belviso I, Carfora A, Montagnani S, Arcucci A: Metabolic Reprogramming of Cancer Associated Fibroblasts: The Slavery of Stromal Fibroblasts. *Biomed Res Int* 2018, 2018: 6075403.
 70. Bakleh MZ, Al Haj Zen A: The Distinct Role of HIF-1 α and HIF-2 α in Hypoxia and Angiogenesis. *Cells* 2025, 14(9): 673.
 71. Sieow JL, Gun SY, Wong SC: The Sweet Surrender: How Myeloid Cell Metabolic Plasticity Shapes the Tumor Microenvironment. *Front Cell Dev Biol* 2018, 6: 168.
 72. Fu Z, Mowday AM, Smaill JB, Hermans IF, Patterson AV: Tumour Hypoxia-Mediated Immunosuppression: Mechanisms and Therapeutic Approaches to Improve Cancer Immunotherapy. *Cells* 2021, 10(5): 1006.
 73. Zhang F, Ma Y, Li D, Wei J, Chen K, Zhang E, Liu G, Chu X, Liu X, Liu W et al: Cancer associated fibroblasts and metabolic reprogramming: unraveling the intricate crosstalk in tumor evolution. *Journal of Hematology & Oncology* 2024, 17(1): 80.
 74. Lee S, Goldfinger LE: RLIP76 regulates HIF-1 activity, VEGF expression and secretion in tumor cells, and secretome transactivation of endothelial cells. *Faseb j* 2014, 28(9): 4158-4168.
 75. Pohl J, Ring A, Korkmaz U, Ehehalt R, Stremmel W: FAT/CD36-mediated long-chain fatty acid uptake in adipocytes requires plasma membrane rafts. *Mol Biol Cell* 2005, 16(1): 24-31.
 76. Ganjoo S, Gupta P, Corbali HI, Nanez S, Riad TS, Duong LK, Barsoumian HB, Masrourpour F, Jiang H, Welsh JW et al: The role of tumor metabolism in modulating T-Cell activity and in optimizing immunotherapy. *Front Immunol* 2023, 14: 1172931.
 77. Wang H, Zhou F, Qin W, Yang Y, Li X, Liu R: Metabolic regulation of myeloid-derived suppressor cells in tumor immune microenvironment: targets and therapeutic strategies. *Theranostics* 2025, 15(6): 2159-2184.
 78. Bailey CM, Liu Y, Liu M, Du X, Devenport M, Zheng P, Liu Y, Wang Y: Targeting HIF-1 α abrogates PD-L1-mediated immune evasion in tumor microenvironment but promotes tolerance in normal tissues. *J Clin Invest* 2022, 132(9): e150846.
 79. Li H, Zhao A, Li M, Shi L, Han Q, Hou Z: Targeting T-cell metabolism to boost immune checkpoint inhibitor therapy. *Front Immunol* 2022, 13: 1046755.
 80. Kierans SJ, Taylor CT: Regulation of glycolysis by the hypoxia-inducible factor (HIF): implications for cellular physiology. *J Physiol* 2021, 599(1): 23-37.
 81. Schiliro C, Firestein BL: Mechanisms of Metabolic Reprogramming in Cancer Cells Supporting Enhanced Growth and Proliferation. *Cells* 2021, 10(5): 1056.
 82. Fricke I, Mirza N, Dupont J, Lockhart C, Jackson A, Lee J-H, Sosman JA, Gabrilovich DI: Vascular Endothelial Growth Factor-Trap Overcomes Defects in Dendritic Cell Differentiation but Does Not Improve Antigen-Specific Immune Responses. *Clinical Cancer Research* 2007, 13(16): 4840-4848.
 83. Albano D, Bosio G, Tomasini D, Bonù M, Giubbini R, Bertagna F: Metabolic behavior and prognostic role of pretreatment 18F-FDG PET/CT in gist. *Asia Pac J Clin Oncol* 2020, 16(5): e207-e215.
 84. Wang Z, Jiang Q, Dong C: Metabolic reprogramming in triple-negative breast cancer. *Cancer Biol Med* 2020, 17(1): 44-59.
 85. Aredo JV, Jamali A, Zhu J, Heater N, Wakelee HA, Vaklavas C, Anagnostou V, Lu J: Liquid Biopsy Approaches for Cancer Characterization, Residual Disease Detection, and Therapy Monitoring. *Am Soc Clin Oncol Educ Book* 2025, 45(3): e481114.
 86. Borger DR, Goyal L, Yau T, Poon RT, Ancukiewicz M, Deshpande V, Christiani DC, Liebman HM, Yang H, Kim H et al: Circulating oncometabolite 2-hydroxyglutarate is a potential surrogate biomarker in patients with isocitrate dehydrogenase-mutant intrahepatic cholangiocarcinoma. *Clin Cancer Res* 2014, 20(7): 1884-1890.
 87. Bobulescu IA, Pop LM, Mani C, Turner K, Rivera C, Khatoon S, Kairamkonda S, Hannan R, Palle K: Renal Lipid Metabolism Abnormalities in Obesity and Clear Cell Renal Cell Carcinoma. *Metabolites* 2021, 11(9): 608.
 88. Vovdenko S, Morozov A, Avraamova S, Alexandrov N, Zharkov N, Kozlov V, Kogan E, Bezrukov E: The role of glucose transporter type 1 (GLUT1) and carbonic anhydrase IX (CAIX) expression by prostate adenocarcinoma tissue in determining disease prognosis and effectiveness of radical treatment. *Urology Herald* 2022, 10(4): 13-20.
 89. Hansen AE, Gutte H, Holst P, Johannesen HH, Rahbek S, Clemmensen AE, Larsen MME, Schøier C, Ardenkjaer-Larsen J, Klausen TL et al: Combined hyperpolarized (13)C-pyruvate MRS and (18)F-FDG PET (hyperPET) estimates of glycolysis in canine cancer patients. *Eur J Radiol* 2018, 103: 6-12.
 90. Richter S, Geldon L, Pang Y, Peitzsch M, Huynh T, Leton R, Viana B, Ercolino T, Mangelis A, Rapizzi E et al: Metabolome-guided genomics to identify pathogenic variants in isocitrate dehydrogenase, fumarate hydratase, and succinate dehydrogenase genes in pheochromocytoma and paraganglioma. *Genet Med* 2019, 21(3): 705-717.
 91. Trotter J, Pantel AR, Teo BK, Escorcia FE, Li T, Pryma DA, Taunk NK: Positron Emission Tomography (PET)/Computed Tomography (CT) Imaging in Radiation Therapy Treatment Planning: A Review of PET Imaging Tracers and Methods to Incorporate PET/CT. *Adv Radiat Oncol* 2023, 8(5): 101212.
 92. Mattoni S, Paccagnella A, Fanti S: The role of fluorine-18-fluorodeoxyglucose-positron emission tomography/computed tomography (18F-FDG-PET/CT) in staging and restaging of patients with uterine sarcomas: a systematic review. *Gynecology and Pelvic Medicine* 2021, 5: 7.
 93. Neumann K, Flavell R, Wilson DM: Exploring Metabolism In Vivo Using Endogenous (11)C Metabolic Tracers. *Semin Nucl Med* 2017, 47(5): 461-473.
 94. Wang ZJ, Ohliger MA, Larson PEZ, Gordon JW, Bok RA, Slater J, Villanueva-Meyer JE, Hess CP, Kurhanewicz J, Vigneron DB: Hyperpolarized (13)C MRI: State of the Art and Future Directions. *Radiology* 2019, 291(2): 273-284.
 95. Jhaveri KS, Elmi A, Hosseini-Nik H, Hedgire S, Evans A, Jewett M, Harisinghani M: Predictive Value of Chemical-Shift MRI in Distinguishing Clear Cell Renal Cell Carcinoma From Non-Clear Cell Renal Cell Carcinoma and Minimal-Fat Angiomyolipoma. *AJR Am J Roentgenol* 2015, 205(1): W79-86.
 96. Yu L, Chen X, Wang L, Chen S: The sweet trap in tumors: aerobic glycolysis and potential targets for therapy. *Oncotarget* 2016, 7(25): 38908-38926.
 97. Roviello G, De Gennaro I, Vascotto I, Venturi G, D'Angelo A, Winchler C, Guarino A, Cacioppo S, Modesti M, Mela MM et

- al: Hypoxia-Inducible Factor in Renal Cell Carcinoma: From Molecular Insights to Targeted Therapies. *Genes (Basel)* 2024, 16(1): 6.
98. Wu H, Fu M, Wu M, Cao Z, Zhang Q, Liu Z: Emerging mechanisms and promising approaches in pancreatic cancer metabolism. *Cell Death & Disease* 2024, 15(8): 553.
 99. Yi M, Jiao D, Qin S, Chu Q, Wu K, Li A: Synergistic effect of immune checkpoint blockade and anti-angiogenesis in cancer treatment. *Mol Cancer* 2019, 18(1): 60.
 100. Qian Y, Yin Y, Zheng X, Liu Z, Wang X: Metabolic regulation of tumor-associated macrophage heterogeneity: insights into the tumor microenvironment and immunotherapeutic opportunities. *Biomark Res* 2024, 12(1): 1.
 101. Kurhanewicz J, Vigneron DB, Ardenkjaer-Larsen JH, Bankson JA, Brindle K, Cunningham CH, Gallagher FA, Keshari KR, Kjaer A, Laustsen C et al: Hyperpolarized (¹³C) MRI: Path to Clinical Translation in Oncology. *Neoplasia* 2019, 21(1): 1-16.
 102. Di Cosimo S, Ferretti G, Papaldo P, Carlini P, Fabi A, Cognetti F: Lonidamine: efficacy and safety in clinical trials for the treatment of solid tumors. *Drugs Today (Barc)* 2003, 39(3): 157-174.
 103. Curry L, Soleimani M: Belzutifan: a novel therapeutic for the management of von Hippel-Lindau disease and beyond. *Future Oncol* 2024, 20(18): 1251-1266.
 104. Lee CH, Motzer R, Enamekhoo H, Matrana M, Percent I, Hsieh JJ, Hussain A, Vaishampayan U, Liu S, McCune S et al: Telaglenastat plus Everolimus in Advanced Renal Cell Carcinoma: A Randomized, Double-Blinded, Placebo-Controlled, Phase II ENTRATA Trial. *Clin Cancer Res* 2022, 28(15): 3248-3255.
 105. Jones SF, Infante JR: Molecular Pathways: Fatty Acid Synthase. *Clin Cancer Res* 2015, 21(24): 5434-5438.
 106. Cheng HJ, Chen NF, Chen WF, Wu ZS, Sun YY, Teng WN, Su FW, Sung CS, Wen ZH: Intrathecal lactate dehydrogenase A inhibitors FX11 and oxamate alleviate chronic constriction injury-induced nociceptive sensitization through neuroinflammation and angiogenesis. *J Headache Pain* 2024, 25(1): 207.
 107. Hasanov E, Jonasch E: MK-6482 as a potential treatment for von Hippel-Lindau disease-associated clear cell renal cell carcinoma. *Expert Opin Investig Drugs* 2021, 30(5): 495-504.
 108. Mondal S, Roy D, Sarkar Bhattacharya S, Jin L, Jung D, Zhang S, Kalogera E, Staub J, Wang Y, Xuyang W et al: Therapeutic targeting of PFKFB3 with a novel glycolytic inhibitor PFK158 promotes lipophagy and chemosensitivity in gynecologic cancers. *Int J Cancer* 2019, 144(1): 178-189.
 109. Basheeruddin M, Qausain S: Hypoxia-Inducible Factor 1-Alpha (HIF-1 α) and Cancer: Mechanisms of Tumor Hypoxia and Therapeutic Targeting. *Cureus* 2024, 16(10): e70700.
 110. Wray R, Mauguen A, Michaud L, Leithner D, Yeh R, Riaz N, Mirtcheva R, Sherman E, Wong R, Humm J et al: Development of (18)F-Fluoromisonidazole Hypoxia PET/CT Diagnostic Interpretation Criteria and Validation of Interreader Reliability, Reproducibility, and Performance. *J Nucl Med* 2024, 65(10): 1526-1532.
 111. Tan SK, Hougén HY, Merchan JR, Gonzalgo ML, Welford SM: Fatty acid metabolism reprogramming in ccRCC: mechanisms and potential targets. *Nat Rev Urol* 2023, 20(1): 48-60.
 112. Zhao Q, Lin X, Wang G: Targeting SREBP-1-Mediated Lipogenesis as Potential Strategies for Cancer. *Front Oncol* 2022, 12: 952371.
 113. Schulte ML, Fu A, Zhao P, Li J, Geng L, Smith ST, Kondo J, Coffey RJ, Johnson MO, Rathmell JC et al: Pharmacological blockade of ASCT2-dependent glutamine transport leads to antitumor efficacy in preclinical models. *Nat Med* 2018, 24(2): 194-202.
 114. Kao TW, Bai GH, Wang TL, Shih IM, Chuang CM, Lo CL, Tsai MC, Chiu LY, Lin CC, Shen YA: Novel cancer treatment paradigm targeting hypoxia-induced factor in conjunction with current therapies to overcome resistance. *J Exp Clin Cancer Res* 2023, 42(1): 171.
 115. Yang WH, Qiu Y, Stamatatos O, Janowitz T, Lukey MJ: Enhancing the Efficacy of Glutamine Metabolism Inhibitors in Cancer Therapy. *Trends Cancer* 2021, 7(8): 790-804.
 116. Rai G, Brimacombe KR, Mott BT, Urban DJ, Hu X, Yang SM, Lee TD, Cheff DM, Kouznetsova J, Benavides GA et al: Discovery and Optimization of Potent, Cell-Active Pyrazole-Based Inhibitors of Lactate Dehydrogenase (LDH). *J Med Chem* 2017, 60(22): 9184-9204.
 117. Dall'Olio FG, Zrafi W, Roelants V, Ambrosini V, Fourquet A, Mitea C, Passiglia F, Bauckneht M, Bonardel G, Conci N et al: Metabolic Tumor Volume Assessed by 18F FDG-PET CT Scan as a Predictive Biomarker for Immune Checkpoint Blockers in Advanced NSCLC and Its Biological Correlates. *Clin Cancer Res* 2025, 31(2): 352-364.
 118. Ahmed N, Escalona R, Leung D, Chan E, Kannourakis G: Tumour microenvironment and metabolic plasticity in cancer and cancer stem cells: Perspectives on metabolic and immune regulatory signatures in chemoresistant ovarian cancer stem cells. *Seminars in Cancer Biology* 2018, 53: 265-281.
 119. Hao ZN, Tan XP, Zhang Q, Li J, Xia R, Ma Z: Lactate and Lactylation: Dual Regulators of T-Cell-Mediated Tumor Immunity and Immunotherapy. *Biomolecules* 2024, 14(12): 1646.
 120. You Y, Lai X, Pan Y, Zheng H, Vera J, Liu S, Deng S, Zhang L: Artificial intelligence in cancer target identification and drug discovery. *Signal Transduction and Targeted Therapy* 2022, 7(1): 156.
 121. Wahi K, Freidman N, Wang Q, Devadason M, Quek LE, Pang A, Lloyd L, Larance M, Zanini F, Harvey K et al: Macropinocytosis mediates resistance to loss of glutamine transport in triple-negative breast cancer. *Embo j* 2024, 43(23): 5857-5882.
 122. Audano M, Pedretti S, Crestani M, Caruso D, De Fabiani E, Mitro N: Mitochondrial dysfunction increases fatty acid β -oxidation and translates into impaired neuroblast maturation. *FEBS Lett* 2019, 593(22): 3173-3189.
 123. Rankin EB, Giaccia AJ: The role of hypoxia-inducible factors in tumorigenesis. *Cell Death Differ* 2008, 15(4): 678-685.
 124. Zhang Y, Shi M, Li M, Qin S, Miao D, Bai Y: Dynamic single-cell metabolomics reveals cell-cell interaction between tumor cells and macrophages. *Nat Commun* 2025, 16(1): 4582.
 125. Li AM, Ye J: Reprogramming of serine, glycine and one-carbon metabolism in cancer. *Biochim Biophys Acta Mol Basis Dis* 2020, 1866(10): 165841.



Copyright © 2025 Ann Urol Oncol. This work is licensed under a Creative Commons Attribution-NonCommercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) License.